

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 7 : A61K 31/22, A61P 3/06, A61K 9/28	A1	(11) International Publication Number: WO 00/53173 (43) International Publication Date: 14 September 2000 (14.09.00)
--	----	---

(21) International Application Number: PCT/US00/02626
(22) International Filing Date: 2 February 2000 (02.02.00)

(30) Priority Data:
60/123,227 8 March 1999 (08.03.99) US
09/264,744 9 March 1999 (09.03.99) US

(71) Applicant (for all designated States except US): MERCK & CO., INC. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): TILLYER, Richard, D. [GB/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US). REIDER, Paul, J. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US). GRABOWSKI, Edward, J., J. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US). XU, Feng [CN/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US). VEGA, Jose, M. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US).

(74) Common Representative: MERCK & CO., INC.; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US).


(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published

With international search report.

With amended claims and statement.

(54) Title: DIHYDROXY OPEN-ACID AND SALTS OF HMG-CO-A REDUCTASE INHIBITORS



(57) Abstract

The instant invention provides methods and pharmaceutical compositions for inhibiting HMG-CoA reductase, as well as for treating and/or reducing the risk for diseases and conditions affected by inhibition of HMG-CoA reductase, comprising orally administering a therapeutically effective amount of a compound selected from a dihydroxy open acid statin and a pharmaceutically acceptable salt or ester thereof in a delayed-release pharmaceutical dosage form to a patient in need of such treatment wherein substantial release of the compound from the dosage form is delayed until after passage of the dosage form through the stomach.

BEST AVAILABLE COPY

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

TITLE OF THE INVENTION

DIHYDROXY OPEN-ACID AND SALTS OF HMG-CO-A REDUCTASE
INHIBITORS

5 FIELD OF THE INVENTION


The instant invention relates to the use of dihydroxy open acid statins and salts and esters thereof, which are inhibitors of 3-hydroxy-3-methylglutaryl coenzyme A (HMG-CoA) reductase, in such a way so as to minimize their in vivo lactonization, and to a particular crystalline hydrated form of the calcium salt of
10 dihydroxy open acid simvastatin referred to herein as compound I.

BACKGROUND OF THE INVENTION


It has been clear for several decades that elevated blood cholesterol is a major risk factor for coronary heart disease (CHD), and many studies have shown that
15 the risk of CHD events can be reduced by lipid-lowering therapy. Prior to 1987, the lipid-lowering armamentarium was limited essentially to a low saturated fat and cholesterol diet, the bile acid sequestrants (cholestyramine and colestipol), nicotinic acid (niacin), the fibrates and probucol. Unfortunately, all of these treatments have limited efficacy or tolerability, or both. With the introduction of lovastatin
20 (MEVACOR®; see US Patent No. 4,231,938), the first inhibitor of HMG-CoA reductase to become available for prescription in 1987, for the first time physicians were able to obtain comparatively large reductions in plasma cholesterol with very few adverse effects.

In addition to the natural fermentation products, mevastatin and
25 lovastatin, there are now a variety of semi-synthetic and totally synthetic HMG-CoA reductase inhibitors, including simvastatin (ZOCOR®; see US Patent No. 4,444,784), pravastatin sodium salt (PRAVACHOL®; see US Patent No. 4,346,227), fluvastatin sodium salt (LESCOL®; see US Patent No. 5,354,772), atorvastatin calcium salt (LIPITOR®; see US Patent No. 5,273,995) and cerivastatin sodium salt (also known
30 as rivastatin; see US Patent No. 5,177,080). The structural formulas of these and additional HMG-CoA reductase inhibitors, are described at page 87 of M. Yalpani, "Cholesterol Lowering Drugs", Chemistry & Industry, pp. 85-89 (5 February 1996). The HMG-CoA reductase inhibitors described above belong to a structural class of compounds which contain a moiety which can exist as either a 3-hydroxy lactone ring
35 or as the corresponding ring opened dihydroxy open-acid, and are often referred to as

"statins." An illustration of the lactone portion of a statin and its corresponding open-acid form is shown below.





Lactone



Dihydroxy Open-Acid

- 5 Salts of the dihydroxy open-acid can be prepared, and in fact, as noted above, several of the marketed statins are administered as the dihydroxy open acid salt forms. Lovastatin and simvastatin are marketed worldwide in their lactonized form. Lovastatin is shown as structural formula II, and simvastatin is shown as structural formula III, below.

10



~~The lactonized forms of the statins are not active inhibitors of HMG-CoA reductase, but the dihydroxy open acid forms are. It is known that condensation~~

- 15 occurs under acidic conditions, that is at about pH 4 or under. Therefore, due to the low gastric pH of the stomach, a statin conventionally administered by oral dosing in its lactone form will remain largely in its lactone form in the stomach. The vast majority of the drug will still be in the lactone form at the time of absorption from the intestine following oral dosing with the lactone. After absorption, the lactone enters
20 the liver and it is in the hepatocytes that the lactone can be metabolized to the active

open acid form, a reaction catalyzed by two hepatic esterases or "lactonases," one which is in the cytosolic and the other in the microsomal fraction. Once in the blood there is an additional plasma esterase which can also hydrolyze the lactone to the open acid. There may be some minimal chemical, i.e., non-enzymatic, hydrolysis that
5 occurs in blood or in the liver; however, at the pH in blood and liver, there should not be any lactonization, i.e., conversion of open acid back to the lactone.

A statin conventionally administered by oral dosing in its dihydroxy open acid form or a pharmaceutically acceptable salt or ester thereof will tend to convert to its lactone form in the acidic environment of the stomach, so that a mixture
10 of the open ring and the corresponding closed ring forms will co-exist there. For example, see M.J. Kaufman, International Journal of Pharmaceutics, 1990, 66(Dec 1), p. 97-106, which provides hydrolysis data that are used to simulate the extent of drug degradation that occurs in acidic gastric fluids following oral administration of several structurally related hypocholesterolemic agents, including simvastatin and lovastatin,
15 and also see A.S. Kearney, et al., Pharmaceutical Research, 1993, 10(10), p. 1461-1465, which describes the interconversion kinetics and equilibrium of CI-981 (atorvastatin in its free acid form). Therefore, even after conventional oral dosing with a dihydroxy open acid statin or a salt or ester thereof, a mixture of the open acid and the corresponding lactone form of the drug could exist by the time of absorption
20 from the intestine.

The preparation of the naturally occurring compound lovastatin and the semi-synthetic analog simvastatin leads to a mixture of the lactone and the open-ring dihydroxy acid forms. Several procedures have been published describing ways to make simvastatin from lovastatin, and most proceed through a lactone ring opening
25 step at some point in the process and sometimes formation of a salt at the resulting carboxy acid, and end with a ring-closing step in order to make the final simvastatin product. For example, U.S. Patent No. 4,820,850 describes a process for making simvastatin which involves opening the lactone ring of lovastatin and forming an alkyl-amide at the resulting carboxy acid, followed by protection of the two hydroxy
30 groups and methylation of the 8' acyl sidechain. After the methylation step, the hydroxy protecting groups are removed, the amide is hydrolyzed to the free acid and an ammonium salt of the free acid is formed, followed by a step to re-lactonize the ring. In U.S. Patent No. 4,444,784, the 8'-acyl sidechain of lovastatin is removed and the lactone ring opened in the first step, followed by re-lactonization of the ring and
35 protection of its hydroxy group. Next, the 8' position is acylated to introduce the

simvastatin sidechain and a deprotection step is performed to obtain the simvastatin final product. In another process disclosed in U.S. Patent No. 4,582,915, the potassium salt of the ring opened form of lovastatin is methylated at the 8' acyl sidechain, the free acid is then re-generated, and the dihydroxy open acid moiety is re-lactonized.

Since becoming available, millions of doses of simvastatin have been administered and these drugs have developed an excellent safety record. However, as noted in the Physician's Desk Reference (PDR), in rare instances myopathy has been associated with the use of all statins, including simvastatin. The mechanism for statin-related myopathy is currently poorly understood. It is also known that many drugs, including certain statins such as simvastatin, are metabolised in the liver and intestine by the cytochrome P450 3A4 (CYP 3A4) enzyme system. As also noted in the PDR, there are adverse drug interaction concerns if a potent inhibitor of CYP3A4, such as itraconazole, and a CYP3A4-metabolized statin are used together, and some cases of myopathy were found to have occurred in patients taking such a drug combination. Simvastatin has been administered to over 20 million patients worldwide in the past 11 years and has been demonstrated to be remarkably safe. However, the very low risk of myopathy is substantially increased when simvastatin is given together with potent inhibitors of CYP3A4. While the overall safety record for simvastatin is exceptional, it would be desirable to further optimize its safety profile by reducing the potential for drug interactions with inhibitors of CYP3A4. It would also be desirable to further reduce the already low rate of occurrence of myopathy associated with the use of all statins. Statins are among the most widely used drugs in the world, and therefore the benefit of any further optimization of their safety profile would be significant.

SUMMARY OF THE INVENTION

One object of this invention is to minimize or eliminate the in vivo lactonization of a dihydroxy open-acid statin. For oral administration, the dihydroxy open-acid statin or a pharmaceutically acceptable salt or ester thereof is to be administered so as to minimize formation of lactonized statin and thereby minimize the amount of lactonized statin that is absorbed from the intestine while maximizing the amount of dihydroxy open-acid statin that is absorbed from the intestine.

Accordingly, this invention involves a method of inhibiting HMG-CoA reductase with an effective inhibitory amount of an orally dosed statin comprising

delivering at least 90% of the dosed statin in its dihydroxy open acid form to the intestinal mucosa of a patient in need of such treatment.

The instant invention further provides a method for inhibiting HMG-CoA reductase, as well as for treating and/or reducing the risk for diseases and conditions affected by inhibition of HMG-CoA reductase, comprising orally administering a therapeutically effective amount of a compound selected from a dihydroxy open acid statin and a pharmaceutically acceptable salt or ester thereof in a delayed-release pharmaceutical dosage form to a patient in need of such treatment wherein substantial release of the compound from the dosage form is delayed until after passage of the dosage form through the stomach.

One embodiment of the first object is to provide the above-described method wherein the delayed-release pharmaceutical dosage form is a gel extrusion module (GEM) drug delivery device.

Another embodiment of the first object is to provide the above-described method wherein the delayed-release pharmaceutical dosage form is an enterically coated pharmaceutical dosage form. An additional aspect of this embodiment is to provide the above-described method wherein the delayed-release pharmaceutical dosage form is selected from an enterically coated rapid-release pharmaceutical dosage form and an enterically coated time controlled-release pharmaceutical dosage form. Yet another aspect of this embodiment is to provide the above-described method wherein the delayed-release pharmaceutical dosage form is an enterically coated gel extrusion module (GEM) drug delivery device.

The compound used for the above-described object and embodiments may particularly be dihydroxy open acid simvastatin and the pharmaceutically acceptable salts thereof, and more particularly a pharmaceutically acceptable salt thereof.

A second object of the instant invention is to provide novel HMG-CoA reductase inhibitors which are crystalline hydrated forms of the calcium salt of dihydroxy open acid simvastatin, and particularly the compound referred to herein as compound I.

Additional objects are to provide the use of the crystalline hydrated forms of the calcium salt of dihydroxy open acid simvastatin, particularly compound I, for inhibiting HMG-CoA reductase, as well as for treating and/or reducing the risk for diseases and conditions affected by inhibition of HMG-CoA reductase, and also to provide pharmaceutical formulations, including conventional rapid-release, delayed-

release and time controlled-release formulations, including the GEM drug delivery device and enterically coated dosage forms, that can be used with the compounds. A further object is to provide a process for making compound I. Additional objects will be evident from the following detailed description.

5

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is a thermogravimetry (TG) weight loss curve for compound I obtained under a nitrogen flow at a heating rate of 10°C/minute.

Figure 2 is a differential scanning calorimetry (DSC) curve for
10 compound I obtained under a nitrogen flow bubbled through 16.0°C water at a heating rate of 10°C/minute in an open cup.

Figure 3 is an x-ray powder diffraction (XRPD) pattern for compound I. The XRPD pattern was obtained using CuK α radiation. The ordinate or Y-axis is x-ray intensity (cpm) and the abscissa or X-axis is the angle two-theta (2 θ) in degrees.

15 Figure 4 is a solid-state ¹³C nuclear magnetic resonance spectrum for Compound I.


DETAILED DESCRIPTION OF THE INVENTION

Applicants have now discovered that dihydroxy open acid statins may
20 be less reliant on CYP3A4 metabolism than their closed ring lactonized counterparts. The instant invention involves methods and pharmaceutical compositions for orally administering open-ring dihydroxy open acid statins and salts and esters thereof, which are HMG-CoA reductase inhibitors, in such a way so as to minimize conversion to their lactonized counterparts. This allows for delivery of a dihydroxy
25 open acid statin without its lactone counterpart directly to the absorptive mucosa of the small intestine thus allowing for absorption of the open acid statin into the portal circulation, penetration by active open acid statin into hepatocytes to achieve enhanced efficacy, and systemic exposure consisting of open acid moieties. More particularly, delayed-release of an orally administered dihydroxy open acid statin or a
30 pharmaceutically acceptable salt or ester thereof, for example dihydroxy open acid simvastatin or a salt thereof, until after passage through the stomach reduces the amount of lactone formed and absorbed in the body. Maintaining the statin in its open acid form in the body thereby reduces the potential for drug interactions between statins whose metabolism is CYP3A4-mediated and other active agents which inhibit
35 the CYP3A4 enzymatic pathway, and also can provide enhanced efficacy. Moreover,


maintaining the statin in its open acid form in the body may have additional clinical benefits for all statins, even for those statins that are not significantly metabolized by the CYP3A4 enzymatic pathway.

- In addition, a novel crystalline hydrated form of the calcium salt of dihydroxy open acid simvastatin has now been discovered to be a pharmaceutically suitable salt form having desirable physical properties for formulation into an anti-hypercholesterolemic drug composition.

- The term "statin(s)" as used herein is intended to be defined as inhibitors of HMG-CoA reductase which belong to a structural class of compounds that contain a moiety which can exist as either a 3-hydroxy lactone ring or as the corresponding ring opened dihydroxy open acid, wherein the lactone portion of the statin and its corresponding dihydroxy open-acid form is shown below.



Lactone



Dihydroxy Open-Acid

- All hydrates, solvates and polymorphic crystalline forms of HMG-CoA reductase inhibitors having the above-described lactone/dihydroxy open acid moiety are included within the scope of the term "statin(s)". Pharmaceutically acceptable salts and esters of the dihydroxy open-acid statins are included within the scope of the term "statin(s)".

- Statins inhibit HMG-CoA reductase in their dihydroxy open acid form. Compounds which have inhibitory activity for HMG-CoA reductase can be readily identified by using assays well-known in the art. For example, see the assays described or cited in U.S. Patent 4,231,938 at col. 6, and WO 84/02131 at pp. 30-33.

- The term "dihydroxy open acid statin(s)" is intended to be defined as statins which contain the dihydroxy open acid moiety, including pharmaceutically acceptable salts and esters thereof. The phrases "dihydroxy open acid statin(s)" and "dihydroxy open acid statin(s) and the pharmaceutically acceptable salts and esters thereof" are used interchangeably herein and are both intended to encompass the open acid and salt and ester forms of the open acid of the statin, unless otherwise indicated.

All hydrates, solvates and polymorphic crystalline forms are encompassed within the the scope of the term "dihydroxy open acid statin(s)."

In the broadest embodiment, any dihydroxy open acid statin or a pharmaceutically acceptable salt or ester thereof may be used with the present invention. Examples of dihydroxy open acid statins that may be used with the present invention include but are not limited to the dihydroxy open acid forms and pharmaceutically acceptable salts and esters thereof of: lovastatin (see US Patent No. 4,342,767); simvastatin (see US Patent No. 4,444,784), pravastatin, particularly the sodium salt thereof; fluvastatin particularly the sodium salt thereof; atorvastatin, particularly the calcium salt thereof; cerivastatin, particularly the sodium salt thereof, nisvastatin also referred to as NK-104 (see PCT international publication number WO 97/23200) and ZD-4522 (see US Patent No. 5,260,440, and Drugs of the Future, 1999, 24(5), pp. 511-513).

In a narrower embodiment, any dihydroxy open acid statin or a pharmaceutically acceptable salt or ester thereof may be used with the present invention, such as those listed above, provided the statin is not pravastatin or fluvastatin. In a class of this embodiment, the open acid statin includes dihydroxy open acid lovastatin, simvastatin, atorvastatin, cerivastatin, nisvastatin and pharmaceutically acceptable salts thereof. Pharmaceutically acceptable salts of dihydroxy open acid simvastatin, particularly the ammonium and calcium salt forms thereof, are preferred for use in the methods and compositions of this invention. More particularly, the calcium salt of dihydroxy open acid simvastatin includes the crystalline hydrated forms of the calcium salt of dihydroxy open acid simvastatin, and more particularly the hydrated calcium salt of dihydroxy open acid simvastatin referred to herein as compound I.

Herein, the term "pharmaceutically acceptable salts" shall mean non-toxic salts of the compounds employed in this invention which are generally prepared by reacting the free acid with a suitable organic or inorganic base, particularly those formed from cations such as sodium, potassium, aluminum, calcium, lithium, magnesium, zinc and tetramethylammonium, as well as those salts formed from amines such as ammonia, ethylenediamine, N-methylglucamine, lysine, arginine, ornithine, choline, N,N'-dibenzylethylenediamine, chloroprocaine, diethanolamine, procaine, N-benzylphenethylamine, 1-p-chlorobenzyl-2-pyrrolidine-1'-yl-methylbenzimidazole, diethylamine, piperazine, morpholine, 2,4,4-trimethyl-2-pentamine and tris(hydroxymethyl)aminomethane. Pharmaceutically acceptable

esters at the carboxylic acid group can be made by treating a dihydroxy open acid statin with an alcohol. Examples of pharmaceutically acceptable esters of dihydroxy open acid statins include, but are not limited to, -C₁₋₄ alkyl and -C₁₋₄ alkyl substituted with phenyl-, dimethylamino-, and acetylamino. "C₁₋₄ alkyl" herein

- 5 includes straight or branched aliphatic chains containing from 1 to 4 carbon atoms, for example methyl, ethyl, n-propyl, n-butyl, *iso*-propyl, *sec*-butyl and *tert*-butyl.

The instant invention involves methods and pharmaceutical compositions for orally administering open-ring dihydroxy open acid statins and salts and esters thereof, which are HMG-CoA reductase inhibitors, in such a way so as to minimize conversion to their lactonized counterparts. This allows for delivery of a dihydroxy open acid statin without its lactone counterpart directly to the absorptive mucosa of the small intestine thus allowing for absorption of the open acid statin into the portal circulation, penetration by active open acid statin into hepatocytes to achieve enhanced efficacy, and systemic exposure consisting of open acid moieties.

10 More particularly, delayed-release of an orally administered dihydroxy open acid statin or a pharmaceutically acceptable salt or ester thereof, for example dihydroxy open acid simvastatin or a salt thereof, until after passage through the stomach reduces the amount of lactone formed and absorbed in the body. Maintaining the statin in its open acid form in the body thereby reduces the potential for drug

- 20 interactions between statins whose metabolism is CYP3A4-mediated and other active agents which inhibit the CYP3A4 enzymatic pathway, and also can provide enhanced efficacy. Administering a statin in its open acid form in such a way so as to minimize conversion to its lactonized counterpart, for example by using an oral delayed release dosage form, should reduce the potential for drug interaction compared to the
- 25 conventional administration of an open acid statin or its lactonized counterpart, for example by using an oral rapid release dosage form.

An object of this invention is to provide methods for reducing the amount of lactonized statin formed and absorbed in the body after oral administration of a dihydroxy open acid statin in order to achieve enhanced clinical benefits. A way to achieve this is to administer the dihydroxy open acid statin in a delayed-release pharmaceutical dosage form. A delayed-release pharmaceutical dosage form as defined herein is an orally administerable pharmaceutical dosage form or device that does not release a substantial amount of the active compound, i.e., the dihydroxy open-acid statin, until after the dosage form has passed through the stomach.

30

35 Therefore, substantial release of the active compound would initially occur after entry

into the duodenum. By "substantial release," it is intended that 90% or more by weight of the active compound in the delayed-release dosage form is released after entry into the duodenum, and that 10% or less by weight of the active compound in the delayed-release dosage form is released in the stomach, i.e., the geometric mean ratio of the plasma AUC (area under the curve) of active vs. total HMG-CoA reductase inhibitory activity will be greater than or equal to 90%. Particularly, the amount of active compound released in the stomach before entry into the duodenum is 5% or less by weight, i.e., the geometric mean ratio of the plasma AUC of active vs. total HMG-CoA reductase inhibitory activity will be greater than or equal to 95%, and more particularly the amount of active compound released in the stomach before entry into the duodenum is 1% or less by weight, i.e., the geometric mean ratio of the plasma AUC of active vs. total HMG-CoA reductase inhibitory activity will be greater than or equal to 99%.

It is to be understood that metabolism of the dihydroxy open acid statins will occur, primarily in the liver, after orally dosing in a delayed-release dosage form. However, since lactonization of the dihydroxy open acid statin would have been substantially avoided by use of a delayed release dosage form, the active and inactive metabolites that are formed will also be in the dihydroxy open acid form. In essence, if the dihydroxy open acid statin is administered in a delayed release dosage form, the internal exposure to lactonized parent compound and also to lactonized active and inactive metabolites will be minimized.

One example of a suitable delayed-release dosage form is a pH-dependent enterically coated dosage form. The enteric coating will not dissolve in the acidic gastric environment, but will dissolve in the higher pH environment of the duodenum. An enterically coated dosage form will therefore not permit release of any significant amount of the active compound from the dosage form in the stomach, but once the enteric coating dissolves in the duodenum, the active compound will be released. Suitable compositions for enteric coatings that can be used with the present invention are known to those of ordinary skill in the pharmaceutical arts; for example, see L. Lachman, *The Theory and Practice of Industrial Pharmacy*, 3rd Ed., H. Liebermann and J. Kanig contributors (Lea & Febiger, 1986). An example of a suitable enteric coating includes but is not limited to SURETERIC WHITE sold by Colorcon which is composed of polyvinyl acetate phthalate, titanium dioxide, talc, colloidal silicon dioxide, triethyl acetate, polyethylene glycol, sodium bicarbonate, purified stearic acid, and sodium alginate. Many other suitable enteric coating

materials are commercially available and are known in the art. Additional coatings employed in the preparation of the dosage form, such as those used to provide an elegant, aesthetically pleasing final product or for other purposes, may be applied before or after, or before and after, application of the enteric coating.

5 Suitable enterically coated pharmaceutical dosage forms for use with this invention include enterically coated conventional rapid-release (also referred to as immediate-release) pharmaceutical dosage forms wherein the drug is relatively rapidly released once the enteric coating is breached, and enterically coated time-
10 controlled release dosage forms such as but not limited to an enterically coated GEM delivery device, described below. Time controlled-release dosage forms are also well known in the pharmaceutical art, and are designed so as to slowly release the active compound in the body over a period of time, for example over a period of from about 6 to 24 hours. Use of an enteric coated time controlled-release dosage form is
15 preferred with more potent dosage amounts of a dihydroxy open acid statin so as to lower the systemic exposure to the active statin. Whether the dosage form is an enterically coated rapid-release or time-controlled release dosage form, the enteric coating will prevent release of any substantial amount of the active compound from the dosage form in the stomach.

 Enterically coated pharmaceutical dosage forms also include but are
20 not limited to those wherein the dosage form or unit is comprised of the dihydroxy open acid statin in a tablet, capsule or the like that is surrounded by an enteric overcoating, and those wherein the dosage form or unit is a tablet, capsule or the like comprised of enterically coated granules of the dihydroxy open acid statin. Where the dosage form is surrounded by an enteric overcoat, the enteric coating may be the
25 outer-most external coating on the dosage form, or there may be one or more additional finish coatings applied over the enteric coat. In a more limited embodiment, when the delayed-release dosage unit contains enterically coated granules of the drug, the drug is selected from the dihydroxy open-acid form of lovastatin and simvastatin and the pharmaceutically acceptable salt and ester forms
30 thereof, and is more preferably a salt of dihydroxy open acid simvastatin, and most preferably the calcium or ammonium salt thereof. In an alternative embodiment, any dihydroxy open acid statin or pharmaceutically acceptable salt or ester thereof may be used with the present invention, such as those described herein, provided that the statin is not dosed in a single pharmaceutical dosage form or unit comprised of enteric

coated granules of the statin and enteric coated or non-enteric coated granules of aspirin.

An example of a delayed-release dosage form that also functions as a time controlled-release dosage form is described in U.S Patent No. 5,366,738, herein
5 incorporated by reference in its entirety. The controlled-release drug delivery device described in U.S Patent No. 5,366,738 is known as a gel extrusion module (GEM) delivery device. The GEM device is a drug delivery device for the controlled in situ production and release of a dispersion containing a beneficial agent such as a pharmaceutical drug comprising:

- 10 (A) a compressed core prepared from an admixture comprising:
- (i) a therapeutically effective amount of the beneficial agent; and
 - (ii) a polymer which upon hydration forms gelatinous microscopic particles; and
- (B) a water insoluble, water impermeable polymeric coating comprising a
15 polymer and a plasticizer, which surrounds and adheres to the core, the coating having a plurality of formed apertures exposing between about 1 and about 75% of the core surface;
- and wherein the release rate of the beneficial agent from the device is a function of the number and size of the apertures.

20 In the GEM device, the polymer inside the compressed core is preferably selected from sodium polyacrylate, carboxypolymethylenes and the pharmaceutically acceptable salts thereof such as a sodium salt, wherein the carboxypolymethylenes are prepared from acrylic acid crosslinked with allylethers of sucrose or pentaerythritol, and more preferably it is selected from
25 carboxypolymethylenes prepared from acrylic acid crosslinked with allylethers of sucrose or pentaerythritol, and the pharmaceutically acceptable salts thereof. Most preferably, CARBOPOL® 974P and pharmaceutically acceptable salts thereof, particularly the sodium salt, is used as the polymer inside the compressed core. In addition, the compressed core may also contain one or more polymer hydration
30 modulating agents, anti-oxidants, lubricants, fillers and excipients. An optional subcoating may be applied to the compressed core prior to application of the water insoluble coating as an aid in the manufacturing process. The subcoating may be comprised of, for example, hydroxypropyl cellulose and hydroxypropylmethylcellulose. Additional coatings may be applied for aesthetic or
35 functional purposes.

The water insoluble, water impermeable polymeric coating is preferably comprised of (1) a polymer selected from polyvinyl chloride, cellulose acetate, cellulose acetate butyrate, ethylcellulose and combinations of these polymers; and (2) a plasticizer selected from diethylphthalate, dibutylsebacate and triethylcitrate. More preferably, the polymeric coating is comprised of cellulose acetate butyrate and triethyl citrate. The GEM device does not function as an osmotic drug delivery device, hence the release function of the device depends on passage of fluids from the external environment of the body to the internal environment of the compressed core through the formed apertures. It is intended that the terms "water insoluble, water impermeable" used to describe the polymeric coating define a coating which is essentially water insoluble and water impermeable, meaning that the polymeric coating allows minimal to no passage of water through the coating from the external environment of the body to the internal environment of the compressed core, except for the fluid passage that occurs through the drilled apertures, during the period of time the drug is being released from the GEM device in the body. Any minimal amount of water that does pass through the water insoluble, water impermeable polymeric coating is insubstantial and does not significantly contribute to the function of the GEM device, i.e. the release rate of the drug through the apertures. Rather the release rate of the drug from the GEM device is primarily a function of the number and size of the apertures on the device.


For an elegant, aesthetically pleasing final product, an outer finish coat may finally be applied to the GEM delivery device containing colorants, waxes, and the like. The GEM device can also be enterically coated, either before or after the application of additional finish coatings. Even without enteric coating, extrusion of the polymer which carries the drug out from inside the compressed core of the GEM device does not occur to a substantial extent in the acidic pH of the stomach, therefore substantial release of the drug should not occur in the stomach. Further details and examples of the GEM delivery device are described in US Patent No. 5,366,738. The compound employed with the GEM device may particularly be a pharmaceutically acceptable salt of dihydroxy open acid simvastatin, and more particularly the ammonium salt of dihydroxy open acid simvastatin.

The term "patient" includes mammals, especially humans, who use the instant active agents for the prevention or treatment of a medical condition. Administering of the drug to the patient includes both self-administration and administration to the patient by another person. The patient may be in need of

treatment for an existing disease or medical condition, or may desire prophylactic treatment to prevent or reduce the risk for diseases and medical conditions affected by inhibition of HMG-CoA reductase.

The term "therapeutically effective amount" is intended to mean that amount of a drug or pharmaceutical agent that will elicit the biological or medical response of a tissue, a system, animal or human that is being sought by a researcher, veterinarian, medical doctor or other clinician. The term "prophylactically effective amount" is intended to mean that amount of a pharmaceutical drug that will prevent or reduce the risk of occurrence of the biological or medical event that is sought to be prevented in a tissue, a system, animal or human by a researcher, veterinarian, medical doctor or other clinician. Particularly, the dosage a patient receives can be selected so as to achieve the amount of LDL (low density lipoprotein) cholesterol lowering desired; the dosage a patient receives may also be titrated over time in order to reach a target LDL level. The dosage regimen utilizing a dihydroxy open acid statin is selected in accordance with a variety of factors including type, species, age, weight, sex and medical condition of the patient; the severity of the condition to be treated; the potency of the compound chosen to be administered; the route of administration; and the renal and hepatic function of the patient. A consideration of these factors is well within the purview of the ordinarily skilled clinician for the purpose of determining the therapeutically effective or prophylactically effective dosage amount needed to prevent, counter, or arrest the progress of the condition.

The novel compounds of this invention are the crystalline hydrated forms of the calcium salt of dihydroxy open acid simvastatin. A particular crystalline hydrated calcium salt of dihydroxy open acid simvastatin is the one having an x-ray powder diffraction (XRPD) pattern obtained using CuK α radiation characterized by reflections at d-spacings of 30.7, 24.6, 15.9, 11.2, 8.58, 7.31, 6.74, 6.06, 5.35, 5.09, 4.93, 4.60, 3.93, 3.84, 3.67, 3.51 and 3.28Å. For convenience, the crystalline hydrated form of the calcium salt of dihydroxy open acid simvastatin having the above-defined XRPD pattern will also be referred to herein as compound I. Compound I can be represented two-dimensionally as a hydrated form of the following structural formula Ia:



In addition to the XRPD pattern described above, compound I of the instant invention is further characterized by the thermogravimetry (TG) curve shown in Figure I. The TG curve for compound I was obtained under a nitrogen flow at a heating rate of 10°C/minute and is characterized by a 6.3% weight loss from ambient room temperature to a stable weight loss plateau at about 175°C. Additional weight losses due to the onset of decomposition of the compound are observed above about 190°C.

Compound I is still further characterized by the differential scanning calorimetry (DSC) curve shown in Figure 2. The DSC curve for compound I was obtained under a nitrogen flow bubbled through 16.0°C water at a heating rate of 10°C/minute in an open cup, and is characterized by three lower temperature endotherms with peak temperatures of 52, 77 and 100°C and associated heats of 48, 90 and 21 J/g, respectively, and two higher temperature endotherms due to decomposition with peak temperatures of 222 and 241°C and associated heats of 163 and 92 J/g.

Compound I of the instant invention is still further characterized by the ¹H nuclear magnetic resonance (NMR) spectral data, ¹³C NMR and mass spectral (MS) data as given in Example 1 herein.

Compound I is also characterized by the solid-state ¹³C nuclear magnetic resonance spectrum shown in Figure 4, which was completed using a Bruker DSX 400WB NMR system operating at 100.6 MHz for ¹³C and 400.1 MHz for ¹H using a Bruker MAS 400WB BL7 double-resonance probe with a spinning module

housing a 7 mm zirconia rotor with Kel-f end caps. The solid-state ^{13}C nuclear magnetic resonance (NMR) spectrum was acquired using cross polarization (CP), magic-angle spinning (MAS), and high-power (~ 59 kHz) decoupling with variable-amplitude cross-polarization and total sideband suppression. Proton and carbon 90° pulse widths were $4.25\ \mu\text{sec}$ with a contact time of $2.0\ \text{msec}$. The sample was spun at $7.0\ \text{kHz}$ and a total of 1024 scans were collected for the spectrum with a recycle delay of 3.0 seconds. A line broadening of $10\ \text{Hz}$ was applied to the spectrum before FT was performed. Chemical shifts are reported on the TMS scale using the carbonyl carbon of glycine ($176.03\ \text{ppm}$) as a secondary reference.

10 In the methods of treatment and prophylaxis described herein, as well as the pharmaceutical compositions and medicaments, a dihydroxy open acid statin or a pharmaceutically acceptable salt or ester thereof is employed. Preferably the compound employed is a pharmaceutically acceptable salt of a dihydroxy open acid statin, more preferably it is a pharmaceutically acceptable salt of dihydroxy open acid
15 simvastatin such as an ammonium salt or calcium salt, and particularly a crystalline hydrated form of the calcium salt of dihydroxy open acid simvastatin such as compound I. All hydrates, solvates and polymorphic crystalline forms of the above-described compounds and their use are encompassed within scope of the instant invention.

20 The instant invention provides methods for inhibiting HMG-CoA reductase, and for treating lipid disorders including hypercholesterolemia, hypertriglyceridemia and combined hyperlipidemia, comprising administering a therapeutically effective amount of a dihydroxy open acid statin to a person in need of such treatment. Further provided are methods for preventing or reducing the risk of
25 developing atherosclerosis, as well as for halting or slowing the progression of atherosclerotic disease once it has become clinically evident, comprising the administration of a prophylactically or therapeutically effective amount, as appropriate, of a dihydroxy open acid statin to a mammal who is at risk of developing atherosclerosis or who already has atherosclerotic disease.

30 Atherosclerosis encompasses vascular diseases and conditions that are recognized and understood by physicians practicing in the relevant fields of medicine. Atherosclerotic cardiovascular disease including restenosis following revascularization procedures, coronary heart disease (also known as coronary artery disease or ischemic heart disease), cerebrovascular disease including multi-infarct

dementia, and peripheral vessel disease including erectile dysfunction are all clinical manifestations of atherosclerosis and are therefore encompassed by the terms "atherosclerosis" and "atherosclerotic disease."

5 A dihydroxy open acid statin may be administered to prevent or reduce the risk of occurrence, or recurrence where the potential exists, of a coronary heart disease event, a cerebrovascular event, and/or intermittent claudication. Coronary heart disease events are intended to include CHD death, myocardial infarction (i.e., a heart attack), and coronary revascularization procedures. Cerebrovascular events are intended to include ischemic or hemorrhagic stroke (also known as cerebrovascular
10 accidents) and transient ischemic attacks. Intermittent claudication is a clinical manifestation of peripheral vessel disease. The term "atherosclerotic disease event" as used herein is intended to encompass coronary heart disease events, cerebrovascular events, and intermittent claudication. It is intended that persons who have previously experienced one or more non-fatal atherosclerotic disease events are those for whom
15 the potential for recurrence of such an event exists.

Accordingly, the instant invention also provides a method for preventing or reducing the risk of a first or subsequent occurrence of an atherosclerotic disease event comprising the administration of a prophylactically effective amount of a dihydroxy open acid statin to a patient at risk for such an event.

20 The patient may or may not have atherosclerotic disease at the time of administration, or may be at risk for developing it.

The instant invention also provides a method for preventing and/or treating inflammatory diseases or disorders alone or in conjunction with the treatment of conditions described above, comprising the administration of a dihydroxy open-
25 acid statin to a patient in need of such treatment. This includes, for example, the treatment of inflammatory conditions susceptible to treatment with a non-steroidal anti-inflammatory agent, arthritis including rheumatoid arthritis, and degenerative joint diseases (osteoarthritis), dementia, Alzheimer's disease, multiple sclerosis, inflammatory bowel disease, asthma, psoriasis, systemic lupus erythematosus,
30 vasculitis, gout, adrenoleukodystrophy, diabetic retinopathy, nephropathy and diabetes mellitus type II.

Persons to be treated with the instant therapy include those at risk of developing atherosclerotic disease and of having an atherosclerotic disease event. Standard atherosclerotic disease risk factors are known to the average physician
35 practicing in the relevant fields of medicine. Such known risk factors include but are

not limited to hypertension, smoking, diabetes, low levels of high density lipoprotein (HDL) cholesterol, and a family history of atherosclerotic cardiovascular disease. Published guidelines for determining those who are at risk of developing atherosclerotic disease can be found in: National Cholesterol Education Program, 5 Second report of the Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel II), National Institute of Health, National Heart Lung and Blood Institute, NIH Publication No. 93-3095, September 1993; abbreviated version: Expert Panel on Detection, Evaluation, and Treatment of 10 High Blood Cholesterol in Adults, Summary of the second report of the national cholesterol education program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel II), JAMA, 1993, 269, pp. 3015-23. People who are identified as having one or more of the above-noted risk factors are intended to be included in the group of people considered at risk for developing atherosclerotic disease. People identified as having one or more 15 of the above-noted risk factors, as well as people who already have atherosclerosis, are intended to be included within the group of people considered to be at risk for having an atherosclerotic disease event.

The oral dosage amount of the dihydroxy open acid statin, particularly a salt of a dihydroxy open acid statin such as simvastatin, and more particularly the 20 ammonium salt or a crystalline form of the calcium salt of dihydroxy open acid simvastatin such as compound I, is from about 1 to 200 mg/day, and more preferably from about 5 to 160 mg/day. However, dosage amounts will vary depending on factors as noted above, including the potency of the particular compound. Although the active drug of the present invention may be administered in divided doses, for 25 example from one to four times daily, a single daily dose of the active drug is preferred. As examples, the daily dosage amount may be selected from, but not limited to, 5 mg, 10 mg, 15 mg, 20 mg, 25 mg, 40 mg, 50 mg, 75 mg, 80 mg, 100 mg, 150 mg and 160 mg.

The active drug employed in the instant therapy can be administered in 30 such oral forms as tablets, capsules, pills, powders, granules, elixirs, tinctures, suspensions, syrups, and emulsions. Oral formulations are preferred.

For the crystalline hydrated forms of the calcium salt of dihydroxy open acid simvastatin, for example compound I, administration of the active drug can be via any pharmaceutically acceptable route and in any pharmaceutically acceptable 35 dosage form. This includes the use of oral conventional rapid-release, time

controlled-release and delayed-release (such as described above) pharmaceutical dosage forms. An oral delayed-release dosage formulation of the instant drug is preferred, and particularly an enteric overcoat surrounding a rapid-release dosage unit, or the GEM controlled-release drug delivery device with an enteric overcoat

5 surrounding the dosage unit, and most particularly an enteric overcoat surrounding a rapid-release dosage unit. Additional suitable pharmaceutical compositions for use with the present invention are known to those of ordinary skill in the pharmaceutical arts; for example, see Remington's Pharmaceutical Sciences, Mack Publishing Co., Easton, PA.

10 In the methods of the present invention, the active drug is typically administered in admixture with suitable pharmaceutical diluents, excipients or carriers (collectively referred to herein as "carrier" materials) suitably selected with respect to the intended form of administration, that is, oral tablets, capsules, elixirs, syrups and the like, and consistent with conventional pharmaceutical practices.

15 For instance, for oral administration in the form of a tablet or capsule, the active drug component can be combined with a non-toxic, pharmaceutically acceptable, inert carrier such as lactose, starch, sucrose, glucose, modified sugars, modified starches, methyl cellulose and its derivatives, dicalcium phosphate, calcium sulfate, mannitol, sorbitol and other reducing and non-reducing sugars, magnesium

20 stearate, steric acid, sodium stearyl fumarate, glyceryl behenate, calcium stearate and the like. For oral administration in liquid form, the drug components can be combined with non-toxic, pharmaceutically acceptable inert carrier such as ethanol, glycerol, water and the like. Moreover, when desired or necessary, suitable binders, lubricants, disintegrating agents and coloring and flavoring agents can also be

25 incorporated into the mixture. Stabilizing agents such as antioxidants, for example butylated hydroxyanisole (BHA), 2,6-di-tert-butyl-4-methylphenol (BHT), propyl gallate, sodium ascorbate, citric acid, calcium metabisulphite, hydroquinone, and 7-hydroxycoumarin, particularly BHA, propyl gallate and combinations thereof, can also be added to stabilize the dosage forms; the use of at least one stabilizing agent is

30 preferred with the instant composition. Preferably an antioxidant is employed with dihydroxy open acid simvastatin or a salt thereof, and particularly compound I. Other suitable components include gelatin, sweeteners, natural and synthetic gums such as acacia, tragacanth or alginates, carboxymethylcellulose, polyethylene glycol, waxes and the like.

The active drug can also be administered in the form of liposome delivery systems, such as small unilamellar vesicles, large unilamellar vesicles and multilamellar vesicles. Liposomes can be formed from a variety of phospholipids, such as cholesterol, stearylamine or phosphatidylcholines.

- 5 Active drug may also be delivered by the use of monoclonal antibodies as individual carriers to which the compound molecules are coupled. Active drug may also be coupled with soluble polymers as targetable drug carriers. Such polymers can include polyvinyl-pyrrolidone, pyran copolymer, polyhydroxy-propyl-methacrylamide-phenol, polyhydroxy-ethyl-aspartamide-phenol, or
10 polyethyleneoxide-polylysine substituted with palmitoyl residues. Furthermore, active drug may be coupled to a class of biodegradable polymers useful in achieving controlled release of a drug, for example, polylactic acid, polyglycolic acid, copolymers of polylactic and polyglycolic acid, polyepsilon caprolactone, polyhydroxy butyric acid, polyorthoesters, polyacetals, polydihydropyrans,
15 polycyanoacrylates and cross linked or amphipathic block copolymers of hydrogels.

- The instant invention also encompasses a process for preparing a pharmaceutical composition comprising combining a dihydroxy open acid statin with a pharmaceutically acceptable carrier. Also encompassed is the pharmaceutical composition which is made by combining a dihydroxy open acid statin with a
20 pharmaceutically acceptable carrier.

- In a broad embodiment, any suitable additional active agent or agents may be used in combination with the dihydroxy open acid statin in a single dosage formulation, or may be administered to the patient in a separate dosage formulation, which allows for concurrent or sequential administration of the active agents. One or
25 more additional active agents may be administered with a dihydroxy open acid statin. The additional active agent or agents can be lipid lowering compounds or agents having other pharmaceutical activities, or agents that have both lipid-lowering effects and other pharmaceutical activities. Examples of additional active agents which may be employed include but are not limited to HMG-CoA synthase inhibitors; squalene
30 epoxidase inhibitors; squalene synthetase inhibitors (also known as squalene synthase inhibitors), acyl-coenzyme A: cholesterol acyltransferase (ACAT) inhibitors including selective inhibitors of ACAT-1 or ACAT-2 as well as dual inhibitors of ACAT-1 and -2; microsomal triglyceride transfer protein (MTP) inhibitors; probucol; niacin; cholesterol absorption inhibitors such as SCH-58235, which is described in U.S.
35 Patent No.'s 5,767,115 and 5,846,966; bile acid sequestrants; LDL (low density

lipoprotein) receptor inducers; platelet aggregation inhibitors, for example glycoprotein IIb/IIIa fibrinogen receptor antagonists and aspirin; human peroxisome proliferator activated receptor gamma (PPAR γ) agonists including the compounds commonly referred to as glitazones for example troglitazone, pioglitazone and rosiglitazone and, including those compounds included within the structural class known as thiazolidinediones as well as those PPAR γ agonists outside the thiazolidinedione structural class; PPAR α agonists such as clofibrate, fenofibrate including micronized fenofibrate, and gemfibrozil; PPAR dual α/γ agonists; vitamin B₆ (also known as pyridoxine) and the pharmaceutically acceptable salts thereof such as the HCl salt; vitamin B₁₂ (also known as cyanocobalamin); folic acid or a pharmaceutically acceptable salt or ester thereof such as the sodium salt and the methylglucamine salt; anti-oxidant vitamins such as vitamin C and E and beta carotene; beta-blockers; angiotensin II antagonists such as losartan; angiotensin converting enzyme inhibitors such as enalapril and captopril; calcium channel blockers such as nifedipine and diltiazam; endothelial antagonists; agents that enhance ABC1 gene expression; FXR and LXR ligands including both inhibitors and agonists; bisphosphonate compounds such as alendronate sodium; and cyclooxygenase-2 inhibitors such as rofecoxib and celecoxib. Additionally, the dihydroxy open acid statins of this invention, for example compound I, may be used in combination with anti-retroviral therapy in AIDS infected patients to treat lipid abnormalities associated with such treatment, for example but not limited to their use in combination with HIV protease inhibitors such as indinavir, nelfinavir, ritonavir and saquinavir.

A therapeutically or prophylactically effective amount, as appropriate, of a crystalline hydrated form of the calcium salt of dihydroxy open acid simvastatin, for example compound I, can be used for the preparation of a medicament useful for inhibiting HMG-CoA reductase, as well as for treating and/or reducing the risk for diseases and conditions affected by inhibition of HMG-CoA reductase, such as treating lipid disorders, preventing or reducing the risk of developing atherosclerotic disease, halting or slowing the progression of atherosclerotic disease once it has become clinically manifest, and preventing or reducing the risk of a first or subsequent occurrence of an atherosclerotic disease event. For example, the medicament may be comprised of about 1 mg to 200 mg of a crystalline hydrated form of the calcium salt of dihydroxy open acid simvastatin, or more particularly about 5 mg to 160 mg.

A therapeutically or prophylactically effective amount, as appropriate, of a dihydroxy open acid statin can be used for the preparation of an oral medicament adapted for delayed-release, wherein substantial release of the statin after oral administration does not occur until after passage of the medicament through the stomach, or alternately wherein at least 90% of the statin is delivered in its dihydroxy open acid form to the intestinal mucosa of a patient after oral administration. Said oral medicaments are also useful for inhibiting HMG-CoA reductase, as well as for treating and/or reducing the risk for diseases and conditions affected by inhibition of HMG-CoA reductase, as described above.

The medicament comprised of a dihydroxy open acid statin, for example compound I, may also be prepared with one or more additional active agents, such as those described supra.

Simvastatin is a semi-synthetic product which can be made from the natural product lovastatin. Processes for preparing lovastatin and simvastatin are well documented in the published literature. For example, U.S. Patent No. 4,231,938, herein incorporated by reference, describes a fermentation and isolation process for obtaining lovastatin using the microorganism *Aspergillus terreus*. U.S. Patent No.'s 4,444,784, 4,820,850, 4,916,239 and 4,582,915, herein all incorporated by reference, describe methods for making dihydroxy open-acid and lactonized forms of simvastatin.

Compound I of the instant invention can generally be prepared as follows. Simvastatin and its dihydroxy open acid counterpart, including compound I, tend to form oxidative by-products; therefore, to minimize the formation of such by-products, it is preferred that the procedures used to make compound I are performed under an inert atmosphere such as nitrogen. Although compound I can be obtained without using an inert atmosphere, the purity of the desired product will not be optimized.

Hydrolysis of the lactone ring of simvastatin can be accomplished by treating simvastatin with at least one equivalent, and preferably a slight excess of one equivalent, of an aqueous base. If more than a slight excess of base is used, the excess base is preferably neutralized before proceeding to the salt formation step in order to prevent formation of insoluble calcium hydroxide or calcium carbonate by-product. The base employed for the hydrolysis can be an aqueous solution of a metal hydroxide or metal carbonate, for example but not limited to sodium hydroxide, potassium hydroxide, lithium hydroxide, sodium carbonate and potassium carbonate.

- Sodium hydroxide is preferred. The hydrolysis can be performed in water, an aqueous-protic organic solvent mixture, or an aqueous-aprotic organic solvent mixture. Suitable protic organic solvents include but are not limited to methanol (MeOH), ethanol (EtOH), isopropyl alcohol, n-propyl alcohol (propanol). Examples of suitable aprotic organic solvents include but are not limited to acetonitrile, dimethyl sulfoxide (DMF), N,N-dimethylformamide (DMSO), tetrahydrofuran (THF), *tert*-butyl methyl ether (MTBE) and toluene. Particularly, an aqueous ethanol or n-propyl alcohol solvent mixture may be used, and more particularly an aqueous n-propyl alcohol solvent mixture.
- After the hydrolysis reaction is complete, the pH of the reaction mixture is adjusted to about 6 to 11, particularly 6 to 9, and more particularly 7 to 8.5, by addition of an acid. In this pH range, the dihydroxy open acid simvastatin will exist as a metal salt, for example as the sodium salt if the base used in the hydrolysis step is sodium hydroxide or sodium carbonate. Any acid that is capable of forming a soluble calcium salt such as calcium chloride or calcium citrate, is suitable. A soluble calcium salt is intended to be a salt that is soluble in the solvent system employed in the instant process. Preferably an acid such as acetic acid (HOAc) or a mineral acid is employed, particularly HCl.

- The resulting pH-adjusted reaction mixture containing the metal salt of dihydroxy open acid simvastatin is next combined with a solution of about 0.50 to 0.55 equivalents of calcium acetate hydrate $[\text{Ca}(\text{OAc})_2 \cdot x\text{H}_2\text{O}]$ in water or an aqueous-organic solvent mixture, such as aqueous EtOH, MeOH, i-PrOH, n-PrOH, acetonitrile, DMF, DMSO, THF, and particularly aqueous EtOH or aqueous n-propyl alcohol. The pH-adjusted reaction mixture can be added to the calcium acetate hydrate solution, or the calcium acetate hydrate solution can be added to the pH-adjusted reaction mixture. The addition can occur all at once, or optionally it can be performed in portions over time with periods of aging. For example, a small portion, e.g., about one-quarter, of the calcium acetate hydrate solution can be added to the pH-adjusted reaction mixture over a short period of time, for example over about 30 minutes, and then the resulting mixture can be allowed to age for an additional short period of time at room temperature, optionally followed by a further period of aging at a temperature up to about 50°C, for example from about 10°C up to about 50°C, particularly from room temperature up to about 50°C, more particularly from about 30 to 40°C, and most particularly from about 30 to 35°C, after which the remaining calcium acetate hydrate solution can be added in portions over several hours at a

temperature up to about 50°C, for example from about 10°C up to about 50°C, particularly from room temperature up to about 50°C, and more particularly from about 30 to 40°C, and most particularly from about 30 to 35°C. Optionally, the reaction mixture can be seeded with crystalline Compound I.

5 Whether the pH-adjusted reaction mixture and the calcium acetate hydrate solution are combined at once or in portions, the resulting slurry must be aged until turnover of the resulting amorphous calcium salt of dihydroxy open acid simvastatin to the crystalline product is complete, usually for at least several hours. Complete turnover to the crystalline product can be assessed by standard techniques in
10 the art, for example, by examining a sample of the product under a microscope. This aging step can be performed at a temperature up to about 50°C, for example from about 10°C up to about 50°C, particularly from room temperature up to about 50°C, and more particularly from about 30 to 40°C, and most particularly from about 30 to 35°C. During the aging period or periods, the use of lower temperatures will lead to
15 crystallized product; however, it has been found that as the temperature drops, the rate of crystal turnover becomes slower, making the procedure less time-efficient.

 If necessary, the slurry is then allowed to cool to room temperature and is collected by suction filtration. The recovered solid is suction dried under a moist atmosphere (about 30 to 70% relative humidity, particularly 40 to 70%), preferably a
20 moist inert atmosphere such as nitrogen, and particularly at a temperature from about 10 to 40°C, and more particularly 25 to 35°C. The final step of suction filtration in the recovery of compound I should be done under a moist atmosphere, and preferably a moist inert atmosphere in order to minimize oxidative by-products. If an additional step of adding an antioxidant to compound I is performed, as described below, then
25 the final suction filtration is the one performed after combining the antioxidant with compound I.

 As noted above, compound I has a tendency to oxidize upon contact with air, and one way to minimize oxidation is to perform the reaction sequence under an inert atmosphere. Additionally, one or more anti-oxidants such as BHA, BHT,
30 propyl gallate, ascorbic acid, calcium metabisulphite, hydroquinone, nordihydroguaiaracetic acid (NDGA) or 7-hydroxycoumarin can be combined with compound I. This is done by agitating a slurry of compound I with one or more of the antioxidants and recovering the resulting solid by suction filtration.

 Alternatively, the ammonium salt of dihydroxy open acid simvastatin
35 can be used as the starting material to be combined with the calcium acetate hydrate,

thus avoiding the hydrolysis and pH adjustment steps needed when starting with lactonized simvastatin. The other reaction conditions described above, such as solvents, temperatures, etc., can otherwise be employed.

Abbreviations which may appear herein are as follows: MeOH is methanol; EtOH is ethanol; PrOH is propanol; HOAc is acetic acid; MeCN is acetonitrile; DMF is dimethyl sulfoxide; DMSO is N,N-dimethylformamide; Ca(OAc)₂ is calcium acetate; HPLC is high performance liquid chromatography; min. is minutes; h is hour(s); D.I. is de-ionized; NMR is nuclear magnetic resonance; EI MS is electron impact mass spectrum; HR-EI MS is high resolution electron impact mass spectrum; RH is relative humidity. The "seed" used in the examples is Compound I.

EXAMPLE 1

15 Preparation of crystalline hydrated calcium salt of dihydroxy open acid simvastatin (Compound I)

A 22 L four-necked round bottom flask was equipped with a temperature probe, a N₂ inlet, an addition funnel, and an overhead stirrer. 8.0 L of 15% EtOH-H₂O was added and the solution was purged with N₂ for 10 min.

20 Simvastatin (396 g, 0.946 mol) was added, and the slurry was purged with N₂ for 5 min. Then 5N NaOH (198 mL) was added at room temperature. After about 1 hour, the hydrolysis reaction was complete as analyzed by HPLC (> 99.9% conversion). The pH of the reaction solution was adjusted to 7 to 8.5 by addition of 1 N HCl (approx. 65 mL). A solution of Ca(OAc)₂·H₂O (116.6 g, 0.662 mol) in 4.0 L of 60%
25 EtOH-H₂O was purged with nitrogen for 5 min. A 1.0 L portion of this solution was added to solution of the sodium salt over 30 min. The resulting slurry was aged at room temperature for 30 min, and then at 30 to 35° C for 1-2 h.

The rest of Ca(OAc)₂ in EtOH-water was added over approx. 30 min at 30 to 35°C. The slurry was aged at 30 to 35 °C for 5 hours under atmosphere of N₂.

30 The slurry was cooled to room temperature and was collected by suction-filtration. The wet cake was washed with 4 L 30% EtOH-H₂O, 4 L 20% EtOH-H₂O, followed by 6 L x 3 of D.I. water. The solid was suction dried under an atmosphere of moist N₂ (40 to 70% relative humidity) at room temperature for 4 days. Crystalline

hydrated calcium salt of dihydroxy open acid simvastatin was obtained as a white powder.

The calcium salt was delumped with a single pass through a cleaned QUADRO™ COMIL® (Model 197S).

5

HPLC CONDITIONS

Column: YMC Basic 4.6 mm x 25 cm
 Detector: ABS 757 1AU/volt output
 10 Sample solvent: EtOH/CH₃CN/H₂O (1:1:1)
 Column temp: 25 °C (Anal. Dept. runs samples at 5 °C to prevent formation of simvastatin on column).
 Flow rate: 1.5 mL / min.
 Wavelength: 238 and 210 nm

15

Gradient:

Time (min.)	%CH ₃ CN	%H ₂ O (10 mM 2HPO ₄ - KH ₂ PO ₄ , pH = 6.5)
-------------	---------------------	---

0.00	30	70
------	----	----

20.00	45	55
-------	----	----

20

34.00	70	30
-------	----	----

39.00	70	30
-------	----	----

39.50	30	70
-------	----	----

43.00	30	70
-------	----	----

25 Retention time of simvastatin open acid: 17.07 min.

Retention time of simvastatin: 32.90 min.

SPECTRAL DATA

30

¹H NMR (400 MHz, CD₃OD), δ 5.97 (d, J = 9.6 Hz, 1 H), 5.77 (dd, J = 9.6, 5.2 Hz, 1 H), 5.49 (m, 1 H), 5.33 (m, 1 H), 4.17 (m, 1 H), 3.70 (m, 1 H), 2.44 - 2.35 (m, 2 H), 2.42 (dd, J = 15.7, 3.6 Hz, 1 H), 2.31 - 2.27 (m, 1 H), 2.29 (dd, J = 15.7, 8.4 Hz, 1 H), 2.00 (ddd, J = 15.7, 7.6, 2.4 Hz, 1 H), 1.93 (m, 1 H), 1.68 (m, 1 H), 1.61 - 1.55 (m, 2

H), 1.55 (m, 2 H), 1.42 (m, 1 H), 1.32 (m, 1 H), 1.19(m, 1 H), 1.12 (s, 6 H), 1.08 (d, J = 7.2 Hz, 3 H), 0.89 (d, J = 7.2 Hz, 3 H), 0.84 (t, J = 7.6 hz, 3 H) ppm.

¹³C NMR (100.55 MHz, CD₃OD), δ 182.3, 179.3, 134.1, 133.2, 130.3, 129.6, 72.5, 5 69.8, 45.1, 44.4, 44.1, 38.8, 38.3, 36.4, 34.3, 33.9, 32.0, 28.6, 25.9, 25.37, 25.36, 23.7, 14.3, 9.9 ppm

EI MS: m/e: 437 (M+H), 419(M+H-H₂O), 303.

10 HR-EI MS: Calcd. for C₂₅H₃₈O₅ 418.2719; Found: 418.2712.

EXAMPLE 2

Preparation of crystalline hydrated calcium salt of dihydroxy open acid simvastatin
(Compound I) with BHA

15 A 22 L three-necked round bottom flask was equipped with a temperature probe, a N₂ inlet, an additional funnel, and an overhead stirrer. 8.0 L of 15% EtOH-H₂O was added and purged with N₂ for 10 min. Simvastatin (396 g, 0.946 mol) was added, and the slurry purged with N₂ for 5 min. 198 mL of 5N NaOH (0.993 mol, 1.05 equiv.) was then added at room temperature. The hydrolysis reaction
20 is usually complete in 1 h. as analyzed by HPLC (> 99.9% conversion). The pH of the reaction solution was adjusted to 7 to 8.5 by addition of 1 N HCl (about 65 mL).

A solution of Ca(OAc)₂•H₂O(91.7 g, 0.520 mol, 0.55 equiv.) in 4.0 L of 60% EtOH- H₂O was purged with nitrogen for 5 min. 1.0 L of this solution was added to reaction solution over 30 min. The slurry was aged at room temperature for
25 30 min, and then at 30 to 35 °C for 1-2 h. The rest of the Ca(OAc)₂ in EtOH-water was added in portions over 3h hours at 30-35°C. The slurry was allowed to age at 30 to 35 °C for 5 h under an atmosphere of N₂. The slurry was allowed to cool to room temperature and was collected by suction-filtration. The wet cake was washed with 4 L 30% EtOH-H₂O, 4 L 20% EtOH-H₂O, followed by 6 L x 3 of D.I. water. The solid
30 was suction dried under atmosphere of moist N₂ (40 to 70% RH) at room temperature to give 1.7 Kg of wet cake.

The above wet cake was placed in a clean 20 L three necked flask under atmosphere of nitrogen. A solution of BHA (2.603 g, 0.2 wt % equiv) in degassed 15% EtOH H₂O (8.5 L) was added, followed by addition of degassed water

(2.55 L), and the slurry was agitated at room temperature for 1 to 2 h. The solid was collected by suction filtration under atmosphere of moist N₂ with no washing to give 1.49 Kg wet cake. The solid was suction dried under atmosphere of moist N₂ (40 to 70% RH) at room temperature for 4 days. The calcium salt title product was obtained
5 as a white powder (94% yield, 99.4%A at 238 nm, 0.2wt% BHA, KF = 7.3 %wt).

EXAMPLE 3

Preparation of crystalline hydrated calcium salt of dihydroxy open acid simvastatin
10 (Compound I) in aqueous nPrOH: Conventional addition mode

Step 1: Hydrolysis

A 72 L three-necked round bottom flask was equipped with a temperature probe, a N₂ inlet, an additional funnel, and an overhead stirrer. 28.5 L of
15 D.I. H₂O was added and purged with N₂ for 10 min. 1.5 Kg simvastatin was added, followed by 788mL of 5N NaOH in one portion at room temperature. The hydrolysis reaction is usually complete in 2 h. as analyzed by HPLC (> 99.9% conversion). 1.5L of nPrOH was added and the pH of the reaction solution was adjusted to 9.5 to 11.0 by adding 2 N HOAc (about 170 mL).

20

Step 2: Salt Formation

150 g of seed (Compound I) was added to the above solution and the resulting slurry was allowed to warm up to 35 to 40 °C. A solution of Ca(OAc)₂•H₂O (347 g) in 15 L of 20% nPrOH was purged with nitrogen for 5min. and added to slurry
25 over 3 h. The resulting slurry was aged at 35 to 40 °C for 5 h. under an atmosphere of N₂ and then cooled to room temperature. The solid was collected by filtration and was washed with 10% nPrOH-H₂O (15 L x 3).

Step 3: BHA loading

30 The above wet cake (9.1 kg) was transferred into a clean 72 L three necked flask under an atmosphere of nitrogen. A solution of BHA (7.6 g) in degassed 10% nPrOH (45 L) was added and the slurry was agitated at room temperature for 1 h. and filtered under atmosphere of N₂, and then suction dried under atmosphere of moist N₂ (30 to 70% RH) at room temperature for 7 days. 1.78 Kg of Ca salt title

product was obtained as a white powder (94% yield. 99.4%A at 238 nm, 0.2wt% BHA, KF = 6.6 %wt).

EXAMPLE 4

5

Preparation of hydrated crystalline calcium salt of dihydroxy open acid simvastatin in aqueous PrOH: Simultaneous addition mode

The process described in this example allows for keeping half of the batch in the vessel at all times as seed in a semi-continuous process.

10 100 g of simvastatin was hydrolyzed in 1.9 L water as described in Example 3. Then, 100 ml nPrOH was added and the solution pH was adjusted to 9 to 11 with 1 N HOAc. The resulting solution and a solution of $\text{Ca}(\text{OAc})_2 \cdot \text{H}_2\text{O}$ (23.2 g) in 1.0 L of 20% nPrOH were added separately but simultaneously to a suspension of
15 the seed) at 30 to 40 °C over 3 h. After 5 h age at 30 to 40 °C, the slurry was cooled to room temperature, filtered, and loaded with anti-oxidant and dried as described in conventional addition mode process. 95% yield.

EXAMPLE 5

20

Preparation of crystalline hydrated calcium salt of dihydroxy open acid simvastatin (Compound I) in aqueous PrOH: Loading BHA through co-crystallization

Method A: 100 g of simvastatin was hydrolyzed in 1.9 L water as described in Example 3. Then, 100 ml nPrOH was added and the solution pH was
25 adjusted to 9 to 11 with 1 N HOAc. 10wt% seed was added and the slurry was allowed to warm up to 35 to 40 °C. A solution of $\text{Ca}(\text{OAc})_2 \cdot \text{H}_2\text{O}$ (23.2 g) and BHA (540 mg) in 1.0 L of 20% nPrOH were added to the slurry at 35 to 40 °C over 3 h. After 5 h age at 30 to 40 °C, the slurry was cooled to room temperature, filtered and washed with a solution of BHA (0.1g/L) in 10% nPrOH (1L x 3). The wet cake was
30 dried under moist N_2 as described in conventional addition mode process. The final dried Ca salt title product contained 0.2wt% BHA. 95% yield.

Method B: The procedure of Method A was employed with the following change. Instead of adding BHA to a $\text{Ca}(\text{OAc})_2$ solution, the same amount of BHA was added into the pH adjusted solution of hydrolyzed simvastatin at room
35 temperature. The solution was warmed to 35 to 40 °C to dissolve BHA. Then, 10wt%

seed was introduced. The rest of the steps were as described in Method A. The final dried Ca salt title product contained 0.2wt% BHA. 95% yield.

EXAMPLE 6

5

Preparation of crystalline hydrated calcium salt of dihydroxy open acid simvastatin
(Compound I) in aqueous PrOH: Loading BHA/propyl gallate

Starting with 2.0 Kg of simvastatin, the calcium salt of dihydroxy open acid simvastatin was crystallized, isolated, and washed as described in Example 3.

10 The first wet cake was transferred to a clean 100L vessel under atmosphere of N₂. A solution of BHA (9.2g) and propyl gallate (11.2 g) in 50 L 10% nPrOH was added to above vessel. The slurry was aged at room temperature for 1h. The slurry was filtered with no wash. The wet cake was dried under moist N₂. 95% yield. The dried salt was loaded with 0.07wt% propyl gallate and 0.2 wt% BHA.

15

EXAMPLE 7

Preparation of crystalline hydrated calcium salt of dihydroxy open acid simvastatin
(Compound I) in aqueous PrOH: Loading propyl gallate

20 Starting with 2.0 Kg of simvastatin, the calcium salt of dihydroxy open acid simvastatin was crystallized, isolated, and washed as described in Example 3. Then, the wet cake was washed with 10L of a solution of propyl gallate in 10% nPrOH (propyl gallate concentration = 0.224 g/L). Then, 20 L of propyl gallate solution in 10% nPrOH (propyl gallate concentration = 0.224 g/L) was added and the
25 wet cake was mixed in the filtration pot before filtration. The wet cake was dried under moist N₂. 95% yield. The dried salt was loaded with 0.07wt% propyl gallate.

EXAMPLE 8

30 Preparation of crystalline hydrated calcium salt of dihydroxy open acid simvastatin
(Compound I) Loading BHA, BHA/Vitamin E, and Vitamin E in heptane

100 g of simvastatin was hydrolyzed in 1.9 L water as described in Example 3. Then, 100 ml nPrOH was added and the solution pH was adjusted to 9 to
35 11 with 1 N HOAc. 10wt% seed was added and the slurry was allowed to warm up to

35 to 40 °C. A solution of $\text{Ca}(\text{OAc})_2 \cdot \text{H}_2\text{O}$ (23.2 g) in 1.0 L of 20% nPrOH was added to a slurry at 35 to 40 °C over 3 h. After 5 h age at 30 to 40 °C, the slurry was cooled to room temperature. The calcium salt slurry was filtered and washed with 10% nPrOH (500 mL x 1), followed by water (1L x 3). The wet cake (KF = 75 to 80 wt% water) was then washed with 1 L of heptane, to displace most of the water. This wet cake was washed with a solution of BHA or Vitamin E or BHA/Vitamin E (conc. = 1.38 g/L, 800 mL) and dried under moist N_2 .

EXAMPLE 9

10

Preparation of crystalline hydrated calcium salt of dihydroxy open acid simvastatin
(Compound I) in aqueous MeCN

A 7.2 L three-necked round bottom flask was equipped with a temperature probe, a N_2 inlet, an additional funnel, and an overhead stirrer. 2.1 L of D.I. H_2O was added and purged with N_2 for 10 min. 150 g simvastatin was added, followed by 78.8 mL of 5N NaOH in one portion at room temperature. The hydrolysis reaction is usually complete in 2 h. as analyzed by HPLC (> 99.9% conversion). 900 mL of MeCN was added and the pH of the reaction solution was adjusted to 9.5 to 11.0 by adding 2 N HOAc (about 17 mL).

30.0 g crystalline seed was added to above solution and the resulting slurry was allowed to warm up to 30 to 35 °C. A solution of $\text{Ca}(\text{OAc})_2 \cdot \text{H}_2\text{O}$ (34.7 g) in 1.5 L of 30% MeCN was purged with nitrogen for 5 min. and added to reaction slurry over 3 h. The slurry was at 35 to 40 °C for 5 h. under atmosphere of N_2 . The slurry was allowed to cool to room temperature and the solid was collected by filtration. The wet cake was washed with 30% MeCN (1.5 L) and 10% MeCN (1.0 L), and rinsed/washed with a solution of BHA (0.9 g/L) in degassed 10% MeCN (1.0 L x 2). The solid was suction dried under atmosphere of moist N_2 (30 to 70% RH) at room temperature for 5 days. 1.67 Kg of the title compound was obtained as a white powder (88% yield. 99.4%A at 238 nm, 0.2wt% BHA, KF = 6.6 %wt).

30

EXAMPLE 10

Preparation of crystalline hydrated calcium salt of dihydroxy open acid simvastatin
(Compound I) in aqueous MeOH

50 g simvastatin was hydrolyzed in 850 mL water as described in Example 3. Then, 150 ml MeOH was added and the solution pH was adjusted to 7 to 11 with 1N HOAc. 10wt% seed was added and the slurry was allowed to warm up to 30-35 °C. A solution of $\text{Ca}(\text{OAc})_2 \cdot \text{H}_2\text{O}$ (11.6 g) in 500 mL of 30% MeOH was added to the slurry at 30-35 °C over 3 h. After 5 h age at 30-35 °C, the slurry was cooled to room temperature. The dihydroxy open acid simvastatin calcium salt slurry was filtered and washed with 20% MeOH (200 ml) and water (500 ml x 3). The wet cake was dried under moist N_2 . The final dried Ca salt title product was isolated in 96% yield.

10

EXAMPLE 11

Preparation of crystalline hydrated calcium salt of dihydroxy open acid simvastatin
(Compound I) in aqueous i-PrOH, DMF, DMSO

15 50 g simvastatin was hydrolyzed in 850 mL water as described in Example 3. Then, 150 ml i-PrOH was added and the solution pH was adjusted to 7 to 11 with 1 N HOAc. 10wt% seed was added and the slurry was allowed to warm up to 30-35 °C. A solution of $\text{Ca}(\text{OAc})_2 \cdot \text{H}_2\text{O}$ (11.6 g) in 500 mL of 30% i-PrOH was added to the slurry at 30-35 °C over 3 h. After 5 h age at 30-35 °C, the slurry was cooled to
20 room temperature. Ca salt slurry was filtered and washed with 20% ml i-PrOH (200 ml) and with water (500 ml x 3). The wet cake was dried under moist N_2 . The final dried Ca salt title product was isolated in 96% yield.

The same procedure could be applied to prepare Compound I in DMF, DMSO, and similar solvents.

25

EXAMPLE 12

Preparation of crystalline hydrated calcium salt of dihydroxy open acid simvastatin
(Compound I) in water

30 50 g of simvastatin was hydrolyzed in 1.0 L water as described in Example 3. Then, the solution pH was adjusted to 11 with 1 N HOAc. 10wt% seed was added and the slurry was allowed to warm up to 35-40 °C. A solution of $\text{Ca}(\text{OAc})_2 \cdot \text{H}_2\text{O}$ (11.6 g) in 500 mL of water was added to the slurry at 35-40 °C over 5 h. After 10 h age at 35-40 °C, the slurry was cooled to room temperature. Ca salt

slurry was filtered and washed with water (500 ml x 3). The wet cake was dried under moist N₂. The final dried Ca salt title product was isolated in 96% yield.

EXAMPLE 13

5

Preparation of crystalline hydrated calcium salt of dihydroxy open acid simvastatin
(Compound I) from dihydroxy open acid simvastatin ammonium salt

Method A: 50 g of dihydroxy open acid simvastatin ammonium salt was dissolved into 800 ml of 25% nPrOH which was then added dropwise to a
10 solution of Ca(OAc)₂•H₂O (10.7 g) in 75 ml of water at room temperature over 2h. The resulting slurry was aged at 30 to 35 °C 5 h. After cooling to room temperature, the slurry was isolated by filtration. The wet cake was washed with 10% nPrOH (500 ml x 3). The wet cake was loaded with antioxidants and dried under moist N₂ as described above to give the title product.

15

Method B: 50 g of dihydroxy open acid simvastatin ammonium salt was added into a solution of Ca(OAc)₂•H₂O (10.7 g) in 1.5 L of 10% nPrOH in one portion at room temperature. The resulting slurry was aged at 30 to 35 °C 5 h. After cooling to room temperature, the slurry was isolated by filtration. The wet cake was washed with 10% nPrOH (500 ml x 3). The wet cake was loaded with antioxidants

20 and dried under moist N₂ as described above to give the title product.

By using both Methods A and B described as above, the title product could also prepared from ammonium salt in the following aqueous solvents: acetone, MeOH, EtOH, iPrOH, , MeCN, neat water, DMF, DMSO, and similar solvents.

25

EXAMPLE 14

Recrystallization Procedure Using nPrOH-H₂O

Dried Compound I (21 g) was dissolved in 150ml of 40% nPrOH at 35°C and line filtered. This solution was added dropwise to a slurry of 10 wt% seed in
30 480 ml of 4% PrOH at 35 to 40°C over 3 to 5 h. After aging overnight at 35 to 40°C, the slurry was allowed to cool to room temperature. The solid was filtered and washed with 10% nPrOH (200ml x 2). The wet cake was dried under moist N₂. 95% yield.

EXAMPLE 15

Recrystallization Process Using EtOH-H₂O

Method A: 25 g of Compound I was dissolved into 425ml of 95% EtOH at 40 °C and line filtered. The filtered solution was added dropwise to 825ml of water in the presence of 10% wt seed at 30 to 35 °C over 3 to 5 h. The slurry was aged overnight and cooled to 0 to 5 °C before filtration. The wet cake was washed with 250 ml of 30% EtOH and dried under moist N₂ at room temperature. 92% yield.

Method B: 25 g of Compound I was dissolved into 625ml of 95% EtOH at 30 to 40 °C and line filtered. 525 ml of water was added at 30 to 40 °C. After adding 10wt% seed, 825 ml of water was added dropwise at 30 to 40 °C over 3 h. The slurry was aged overnight and cooled to 0 to 5 °C before filtration. The wet cake was washed with 250 ml of 30% EtOH and dried under moist N₂ at room temperature. 92% yield.

EXAMPLE 16

An open, randomized, four-period, crossover study to compare the effect of itraconazole on the single-dose pharmacokinetics of intraduodenally administered dihydroxy open acid simvastatin versus orally administered simvastatin in healthy male subjects

Objectives: (1) To determine the effect of itraconazole, a potent CYP3A inhibitor, on the plasma AUC of active and of total HMG-CoA reductase inhibitory activity following a single intraduodenal dose of a solution containing 5 mg dihydroxy open acid simvastatin; (2) to determine and compare the dose-adjusted plasma AUC of active HMG-CoA reductase inhibitory activity following a single intraduodenal dose of a solution containing 5 mg dihydroxy open acid simvastatin versus a single oral administration of simvastatin 20-mg film coated tablet (FCT); (3) to determine the effect of itraconazole on the plasma AUC's of dihydroxy open acid simvastatin and of simvastatin lactone concentration following a single intraduodenal dose of a solution containing 5 mg dihydroxy open acid simvastatin.

Study Design: This study was designed in an open, four-period crossover randomized fashion. Twelve healthy male subjects received four treatments (A, B, C and D). In Treatment A, subjects received itraconazole 200 mg (2 x 100-mg capsule) for 4 days followed by a single dose of 5-mg dihydroxy open acid simvastatin solution administered intraduodenally on day 4, 1 hour after the fourth daily dose of itraconazole. In Treatment B, subjects were given a single dose of 5-mg dihydroxy open acid simvastatin solution administered intraduodenally on day 1. Intraduodenal administration was accomplished via a nasoduodenal tube placed under fluoroscopic guidance by an experienced gastroenterologist just prior to dosing and removed following the 1-hr postdose measurements. Treatments C and D were similar to those of Treatments A and B, except that orally dosed simvastatin 20-mg conventional film coated tablet was used. The wash out between treatment periods was at least 7-days following a treatment containing itraconazole or at least 3 days following a treatment without itraconazole. Plasma samples were collected at appropriate time intervals for up to 24 hours following simvastatin or dihydroxy open acid simvastatin administration, for analysis of total and active HMG-CoA reductase inhibitory activities as well as for simvastatin and dihydroxy open acid simvastatin concentrations.

20 Analytical Methodology: Plasma concentrations of simvastatin and dihydroxy open acid simvastatin acid were determined simultaneously by an improved liquid chromatography/tandem mass spectrometry (LC/MS/MS) method using lovastatin and dihydroxy open acid lovastatin acid as internal standards. An enzymatic assay method was used to determine plasma concentrations of active and total (active plus potentially active) HMG-CoA reductase inhibitory activity.

Pharmacokinetics: The area under the plasma concentration-time profile from time zero to the last sampling time (AUC_{0-last}) was calculated using linear trapezoidal rule. The apparent elimination rate constant (k) of simvastatin and dihydroxy open acid simvastatin was estimated by least-squares regression analysis of the log-linear portion of the simvastatin and dihydroxy open acid simvastatin concentration-time data, and the apparent elimination half-life ($t_{1/2}$) was calculated as $t_{1/2} = 0.693/k$. All calculations were based on designated sampling times or actual sampling times when they differed from the designated times by more than 10 minutes.

35

Discussion of Results: This was an open, randomized, four-period crossover study in twelve healthy male subjects. The results showed that intraduodenal administration of dihydroxy open acid simvastatin 5-mg solution yielded higher (~4-fold) dose-adjusted plasma AUC of the active HMG-CoA reductase inhibitory activity than oral administration of simvastatin 20-mg tablet (Table 1). Following dihydroxy open acid simvastatin administration, the unchanged dihydroxy open acid simvastatin was the major component (~60%), while simvastatin was a minor component (<10%) contributing to plasma HMG-CoA reductase inhibitory activity. As evident by comparable AUC values for both the total and active inhibitors (see Table 1) as well as low plasma levels of simvastatin in plasma (AUC <10% of dihydroxy open acid simvastatin AUC) (see Table 3), lactonization of either dihydroxy open acid simvastatin or its active metabolites occurred minimally following intraduodenal administration of dihydroxy open acid simvastatin. Pretreatment with itraconazole caused minimal changes (1.3-1.5-fold) in the systemic exposure as measured by AUC and C_{max} of HMG-CoA reductase inhibitory activity (total or active) following administration of dihydroxy open acid simvastatin 5-mg intraduodenally, as compared to that observed following oral administration of simvastatin 20-mg tablet (1.3-3.8-fold) (see Table 2). When measured as the unchanged drug, the effect of itraconazole observed following dihydroxy open acid simvastatin administration was also minimal (1.5-fold) and was much less than the corresponding measure obtained following simvastatin administration (19-fold) (see Table 3). A moderate effect (3-4-fold increase) was noted for the AUC and C_{max} of simvastatin following treatment with itraconazole prior to dihydroxy open acid simvastatin administration (see Table 3). However, apparent t_{1/2} values for dihydroxy open acid simvastatin or simvastatin were essentially unchanged by itraconazole (see Table 3). Overall, these results indicate that the pharmacokinetics of dihydroxy open acid simvastatin is less prone to alteration by itraconazole, a potent CYP3A inhibitor, than that of simvastatin in humans. From these results it appears that dihydroxy open acid simvastatin, although a substrate for CYP3A, is metabolized with a much lower intrinsic clearance than that of simvastatin in human liver microsomes.

TABLE 1

Pharmacokinetic parameters for total and active HMG-CoA reductase inhibitors following administration of 5-mg dihydroxy open acid simvastatin (SVA) intraduodenally or 20-mg simvastatin (SV) tablet orally to 12 healthy male volunteers. Results are means from 12 subjects. Values in parentheses are SD.

	Drug Administered	5 mg SVA (ID)	20 mg SV (FCT)
	N	12	12
AUC (ng eq. hr/ml)	Total	56.0 (28.6)	180.7 (57.2)
	Active	54.1 (28.5)	59.7 (18.1)
	SV	1.02 (0.5)	16.9 (10.5)
	SVA	33.9 (16.7)	7.7 (4.9)
Cmax (ng eq./ml)	Total	6.2 (3.6)	67.1 (27.2)
	Active	5.7 (3.0)	16.1 (4.4)
	SV	0.13 (0.05)	6.84 (4.6)
	SVA	3.78 (1.93)	0.92 (0.58)
Tmax (hr)	Total	3.3 (2.3)	1.2 (0.5)
	Active	3.6 (2.5)	1.4 (0.5)
	SV	4.9 (2.0)	1.1 (0.5)
	SVA	4.1 (2.1)	3.7 (2.1)
t1/2 (hr)	SV	6.7 (1.7) n=8	4.2 (1.9)
	SVA	2.3 (0.5)	3.5 (1.0)

TABLE 2

Pharmacokinetic parameters for total HMG-CoA reductase inhibitor following administration of 5-mg dihydroxy open acid simvastatin (SVA) intraduodenally or 20-mg simvastatin (SV) tablet orally with or without pretreatment with itraconazole (2x100-mg capsule, qd) for 4 days to 12 healthy male volunteers. Results are means from 12 subjects. Values in parentheses are SD.

Treatment	AUC (ng eq.hr/mL)	C _{max} (ng eq/mL)	T _{max} (hr)
Total Inhibitors			
Single 5-mg SVA (ID) + 4 Days of Daily 200-mg Itraconazole	86.1 ± 47	8.2 ± 4.0	4.3 ± 2.2
Single 5-mg SVA (ID)	56.0 ± 28.6	6.2 ± 3.6	3.3 ± 2.3
Geometric Mean Ratio	1.5	1.3	
Single 20-mg SV (CT) + 4 Days of Daily 200-mg Itraconazole	683 ± 232	126 ± 54.8	1.9 ± 0.9
Single 20-mg SV (CT)	180.7 ± 57.2	67.1 ± 27.2	1.2 ± 0.5
Geometric Mean Ratio	3.8	1.9	
Active Inhibitors			
Single 5-mg SVA (ID) + 4 Days of Daily 200-mg Itraconazole	75.0 ± 39.6	7.4 ± 3.9	5.5 ± 1.7
Single 5-mg SVA (ID)	54.1 ± 28.5	5.7 ± 3.0	3.6 ± 2.5
Geometric Mean Ratio	1.3	1.3	
Single 20-mg SV (CT) + 4 Days of Daily 200-mg Itraconazole	195 ± 104	23.0 ± 13.6	3.1 ± 0.9
Single 20-mg SV (CT)	59.7 ± 18.1	16.1 ± 4.4	1.4 ± 0.5
Geometric Mean Ratio	3.1	1.3	

TABLE 3

Pharmacokinetic parameters for simvastatin or dihydroxy open acid simvastatin following administration of 5-mg dihydroxy open acid simvastatin (SVA) intraduodenally or 20-mg simvastatin (SV) tablet orally with or without pretreatment with itraconazole (2x100-mg capsule, qd) for 4 days to 12 healthy male volunteers. Results are means from 12 subjects. Values in parentheses are SD.

Treatment	AUC (ng eq.hr/mL)	C _{max} (ng eq/mL)	t _{1/2} (hr)	T _{max} (hr)
Simvastatin				
Single 5-mg SVA (ID) + 4 Days of Daily 200-mg Itraconazole	5.18 ± 3.15	0.43 ± 0.2	6.7 ± 2.5 (n=11)	5.3 ± 1.9
Single 5-mg SVA (ID)	1.02 ± 0.5	0.13 ± 0.05	6.7 ± 1.7 (n=8)	4.9 ± 2.0
Geometric Mean Ratio	4.4	3.2	1.0	
Single 20-mg SV (CT) + 4 Days of Daily 200-mg Itraconazole	316 ± 142	77.2 ± 50.7	5.2 ± 1.5	1.8 ± 1.1
Single 20-mg SV (CT)	16.9 ± 10.5	6.84 ± 4.6	4.2 ± 1.9	1.1 ± 0.5
Geometric Mean Ratio	19.4	11.4	1.3	
Simvastatin Acid				
Single 5-mg SVA (ID) + 4 Days of Daily 200-mg Itraconazole	56.8 ± 33.5	5.78 ± 3.27	2.8 ± 1.1	4.7 ± 2.4
Single 5-mg SVA (ID)	33.9 ± 16.7	3.78 ± 1.93	2.3 ± 0.5	4.1 ± 2.1
Geometric Mean Ratio	1.5	1.5	1.1	
Single 20-mg SV (CT) + 4 Days of Daily 200-mg Itraconazole	86.2 ± 65.6	9.76 ± 7.38	4.5 ± 1.3	4.0 ± 1.4
Single 20-mg SV (CT)	7.70 ± 4.9	0.92 ± 0.58	3.5 ± 1.0	3.7 ± 2.1
Geometric Mean Ratio	11.0	10.7	1.3	

While the invention has been described and illustrated with reference to certain particular embodiments thereof, those skilled in the art will appreciate that various changes, modifications and substitutions can be made therein without departing from the spirit and scope of the invention. For example, effective dosages
5 other than the particular dosages as set forth herein above may be applicable as a consequence of variations in the responsiveness of the mammal being treated for any of the indications for the active agents used in the instant invention as indicated above. Likewise, the specific pharmacological responses observed may vary according to and depending upon the particular active compound selected or whether
10 there are present pharmaceutical carriers, as well as the type of formulation employed, and such expected variations or differences in the results are contemplated in accordance with the objects and practices of the present invention. It is intended, therefore, that the invention be defined by the scope of the claims which follow and that such claims be interpreted as broadly as is reasonable.

WHAT IS CLAIMED IS:

1. An oral pharmaceutical composition comprising a therapeutically effective amount of a compound selected from a dihydroxy open acid statin and a pharmaceutically acceptable salt or ester thereof and a pharmaceutically acceptable carrier, formulated as a delayed-release dosage form wherein substantial release of the compound from the dosage form after oral administration to a patient is delayed until after passage of the dosage form through the stomach.
2. The pharmaceutical composition of Claim 1 wherein at most 10% by weight of the statin is released into the stomach of a patient after administration.
3. The pharmaceutical composition of Claim 2 wherein at most 5% by weight of the statin is released into the stomach of a patient after administration.
4. The pharmaceutical composition of Claim 3 wherein at most 1% by weight of the statin is released into the stomach of a patient after administration.
5. The pharmaceutical composition of Claim 1 wherein the composition is enterically coated.
6. The pharmaceutical composition of Claim 5 wherein the dosage form is surrounded by an enteric coating.
7. The pharmaceutical composition of Claim 5 wherein the dosage form comprises enterically coated granules of the dihydroxy open acid statin or a pharmaceutically acceptable salt or ester thereof.
8. The pharmaceutical composition of Claim 5 wherein the dosage form comprises enterically coated granules of dihydroxy open acid simvastatin or a pharmaceutically acceptable salt or ester thereof.

9. The pharmaceutical composition of Claim 5 wherein the dosage form comprises enterically coated granules of the dihydroxy open acid statin or a pharmaceutically acceptable salt or ester thereof, provided that the dosage form does not contain granules of aspirin.

5

10. The pharmaceutical composition of Claim 5 wherein the composition is formulated in an enterically coated rapid-release pharmaceutical dosage form.

10

11. The pharmaceutical composition of Claim 5 wherein the composition is formulated in an enterically coated time controlled-release pharmaceutical dosage form.

15

12. The pharmaceutical composition of Claim 1 wherein the composition is formulated in a drug delivery device comprised of:

- (A) a compressed core prepared from an admixture comprising:
- (i) a therapeutically effective amount of the compound; and
 - (ii) a polymer which upon hydration forms gelatinous microscopic particles; and

20

- (B) a water insoluble, water impermeable polymeric coating comprising a polymer and a plasticizer, which surrounds and adheres to the core, the coating having a plurality of formed apertures exposing between about 1 and about 75% of the core surface;

and wherein the release rate of the compound from the device is a function of the number and size of the apertures.

25

13. The pharmaceutical composition of Claim 12 wherein the drug delivery device is additionally comprised of an enteric overcoat exterior to the water insoluble, water impermeable polymeric coating.

30

14. The composition of Claim 1 wherein the statin is selected from the dihydroxy open acid forms of lovastatin, simvastatin, atorvastatin, cerivastatin, nisvastatin and ZD-4522 and the pharmaceutically acceptable salts and esters thereof.

15. The composition of Claim 1 wherein the statin is selected from the dihydroxy open acid form of lovastatin, simvastatin and the pharmaceutically acceptable salts and esters thereof.

5 16. The composition of Claim 1 wherein the statin is selected from the dihydroxy open acid form of simvastatin and the pharmaceutically acceptable salts and esters thereof.

10 17. The composition of Claim 1 wherein the statin is a pharmaceutically acceptable salt of dihydroxy open acid simvastatin.

18. The composition of Claim 1 wherein the statin is a calcium salt of dihydroxy open acid simvastatin.

15 19. The composition of claim 1 wherein the statin is an ammonium salt of dihydroxy open acid simvastatin.

20 20. The composition of Claim 1 wherein the statin is a crystalline hydrated calcium salt of dihydroxy open acid simvastatin having an x-ray powder diffraction pattern obtained using CuK α radiation characterized by reflections at d-spacings of 30.7, 24.6, 15.9, 11.2, 8.58, 7.31, 6.74, 6.06, 5.35, 5.09, 4.93, 4.60, 3.93, 3.84, 3.67, 3.51 and 3.28Å.

25 21. The composition of Claim 5 wherein the statin is selected from the dihydroxy open acid form of lovastatin, simvastatin, atorvastatin, cerivastatin, nisvastatin and ZD-4522 and the pharmaceutically acceptable salts and esters thereof.

30 22. The composition of Claim 5 wherein the statin is selected from the dihydroxy open acid form of lovastatin, simvastatin and the pharmaceutically acceptable salts and esters thereof.

35 23. The composition of Claim 5 wherein the statin is selected from the dihydroxy open acid form of simvastatin and the pharmaceutically acceptable salts and esters thereof.

24. The composition of Claim 5 wherein the statin is a pharmaceutically acceptable salt of simvastatin.

25. The composition of Claim 5 wherein the statin is a calcium salt of dihydroxy open acid simvastatin.

26. The composition of Claim 5 wherein the statin is an ammonium salt of dihydroxy open acid simvastatin.

27. The composition of Claim 5 wherein the statin is a crystalline hydrated calcium salt of dihydroxy open acid simvastatin having an x-ray powder diffraction pattern obtained using CuK α radiation characterized by reflections at d-spacings of 30.7, 24.6, 15.9, 11.2, 8.58, 7.31, 6.74, 6.06, 5.35, 5.09, 4.93, 4.60, 3.93, 3.84, 3.67, 3.51 and 3.28Å.

28. The composition of claim 27 further comprising BHA.

29. The composition of claim 27 further comprising propyl gallate.

30. The composition of claim 27 further comprising BHA and propyl gallate.

31. An oral pharmaceutical composition made by combining a therapeutically effective amount of a compound selected from a dihydroxy open acid statin and a pharmaceutically acceptable salt or ester thereof with a pharmaceutically acceptable carrier in a delayed-release dosage form.

32. A process for preparing an oral pharmaceutical composition comprising combining a compound selected from a dihydroxy open acid statin and a pharmaceutically acceptable salt or ester thereof with a pharmaceutically acceptable carrier in a delayed-release dosage form.

33. The use of a dihydroxy open acid statin or a pharmaceutically acceptable salt or ester thereof for the preparation of an oral medicament adapted for

delayed-release wherein substantial release of the statin after oral administration does not occur until after passage of the medicament through the stomach.

5 34. The use of Claim 33 wherein at most 10% by weight of the statin is released into the stomach of a patient after administration.

35. The use of Claim 33 wherein at most 5% by weight of the statin is released into the stomach of a patient after administration.

10 36. The use of Claim 33 wherein at most 1% by weight of the statin is released into the stomach of a patient after administration.

37. The use of dihydroxy open acid simvastatin or a pharmaceutically acceptable salt or ester thereof to prepare the medicament of claim
15 33.

38. The use of a dihydroxy open acid statin or a pharmaceutically acceptable salt or ester thereof for the preparation of an oral medicament adapted for
~~delayed-release useful for delivering at least 90% by weight of the statin in its~~
20 dihydroxy open acid form to the intestinal mucosa of a patient after oral administration.

39. The use of Claim 38 wherein at least 95% by weight of the statin is delivered in its dihydroxy open acid form to the intestinal mucosa of a patient
25 after oral administration.

40. The use of Claim 38 wherein at least 99% by weight of the statin is delivered in its dihydroxy open acid form to the intestinal mucosa of a patient
after oral administration.

30 41. The use of dihydroxy open acid simvastatin or a pharmaceutically acceptable salt or ester thereof to prepare the medicament of claim 38.

42. The use of a dihydroxy open acid statin or a pharmaceutically acceptable salt or ester thereof for the preparation of an oral medicament adapted for delayed-release useful for inhibiting HMG-CoA reductase in a patient in need of such treatment wherein substantial release of the statin after oral administration is delayed
5 until after passage of the medicament through the stomach.

43. The use of a dihydroxy open acid statin or a pharmaceutically acceptable salt or ester thereof for the preparation of an oral medicament adapted for delayed-release useful for treating hypercholesterolemia in a patient in need of such
10 treatment wherein substantial release of the statin after oral administration is delayed until after passage of the medicament through the stomach.

44. The use of a dihydroxy open acid statin or a pharmaceutically acceptable salt or ester thereof for the preparation of an oral medicament adapted for
15 delayed-release useful for preventing or reducing the risk of developing atherosclerotic disease in a patient at risk of developing atherosclerotic disease wherein substantial release of the statin after oral administration is delayed until after passage of the medicament through the stomach.

20 45. The use of Claim 44 wherein the medicament is useful for preventing or reducing the risk of developing atherosclerotic disease selected from cardiovascular disease, cerebrovascular disease and peripheral vessel disease.

46. The use of Claim 44 wherein the medicament is useful for
25 preventing or reducing the risk of developing coronary heart disease.

47. The use of a dihydroxy open acid statin or a pharmaceutically acceptable salt or ester thereof for the preparation of an oral medicament adapted for delayed-release useful for treating atherosclerotic disease in a patient in need of such
30 treatment wherein substantial release of the statin after oral administration is delayed until after passage of the medicament through the stomach.

48. The use of Claim 47 wherein the medicament is useful for treating atherosclerotic disease selected from cardiovascular disease, cerebrovascular
35 disease and peripheral vessel disease.

49. The use of Claim 47 wherein the medicament is useful for treating coronary heart disease.

5 50. The use of a dihydroxy open acid statin or a pharmaceutically acceptable salt or ester thereof for the preparation of an oral medicament adapted for delayed-release useful preventing or reducing the risk of occurrence or recurrence of an atherosclerotic disease event in a person at risk of having an atherosclerotic disease event wherein substantial release of the statin after oral administration is delayed until
10 after passage of the medicament through the stomach.

51. The use of Claim 50 wherein the medicament is useful for preventing or reducing the risk of occurrence or recurrence of an atherosclerotic disease event in a person who has atherosclerotic disease.

15 52. The use of Claim 50 wherein the medicament is useful for preventing or reducing the risk of occurrence or recurrence of an atherosclerotic disease event in a person who is at risk of developing atherosclerotic disease.

20 53. The use of Claim 50 wherein the medicament is useful for preventing or reducing the risk of occurrence or recurrence of an atherosclerotic disease event selected from a coronary heart disease event, a cerebrovascular event and intermittent claudication.

25 54. The use of Claim 50 wherein the medicament is useful for preventing or reducing the risk of occurrence or recurrence of a coronary heart disease event selected from coronary heart disease death, myocardial infarction and coronary revascularization procedures.

30 55. The use of Claim 50 wherein the medicament is useful for preventing or reducing the risk of occurrence or recurrence of a cerebrovascular event selected from a cerebrovascular accident and a transient ischemic attack.

56. The use of any of Claims 42 to 55 wherein the statin is selected from the dihydroxy open acid forms of lovastatin and simvastatin and the pharmaceutically acceptable salts and esters thereof.
- 5 57. The use of any of Claims 42 to 55 wherein the statin is selected from dihydroxy open acid simvastatin and the pharmaceutically acceptable salts and esters thereof.
58. The use of any of Claims 42 to 55 wherein the statin is a
10 pharmaceutically acceptable salt of dihydroxy open acid simvastatin.
59. The use of any of Claims 42 to 55 wherein the statin is a calcium salt of dihydroxy open acid simvastatin.
- 15 60. The use of any of Claims 42 to 55 wherein the statin is an ammonium salt of dihydroxy open acid simvastatin.
61. The use of any of Claims 42 to 55 wherein the statin is a
crystalline hydrated calcium salt of dihydroxy open acid simvastatin having an x-ray
20 powder diffraction pattern obtained using CuK α radiation characterized by reflections at d-spacings of 30.7, 24.6, 15.9, 11.2, 8.58, 7.31, 6.74, 6.06, 5.35, 5.09, 4.93, 4.60, 3.93, 3.84, 3.67, 3.51 and 3.28Å.

AMENDED CLAIMS

[received by the International Bureau on 10 July 2000 (10.07.00)
original claims 8,14-16,21-23,37,41 and 56-57 cancelled original claims
1-4,7,9,17-20,24-36,38-40,42-44,47,50 and 58-61 amended; new claims 62-64 added;
remaining claims unchanged (8 pages)]

- 5 1. An oral pharmaceutical composition comprising a therapeutically effective amount of a compound selected from dihydroxy open acid simvastatin and the pharmaceutically acceptable salts and esters thereof and a pharmaceutically acceptable carrier, formulated as a delayed-release dosage form wherein substantial release of the compound from the dosage form after oral administration to a patient is delayed until after passage of the dosage form through the stomach.
- 10 2. The pharmaceutical composition of Claim 1 wherein at most 10% by weight of the compound is released into the stomach of the patient after oral administration.
- 15 3. The pharmaceutical composition of Claim 2 wherein at most 5% by weight of the compound is released into the stomach of the patient after oral administration.
- 20 4. The pharmaceutical composition of Claim 3 wherein at most 1% by weight of the compound is released into the stomach of the patient after oral administration.
- 25 5. The pharmaceutical composition of Claim 1 wherein the composition is enterically coated.
6. The pharmaceutical composition of Claim 5 wherein the dosage form is surrounded by an enteric coating.
- 30 7. The pharmaceutical composition of Claim 5 wherein the dosage form comprises enterically coated granules of the compound.

9. The pharmaceutical composition of Claim 5 wherein the dosage form comprises enterically coated granules of the compound, provided that the dosage form does not contain granules of aspirin.

5 10. The pharmaceutical composition of Claim 5 wherein the composition is formulated in an enterically coated rapid-release pharmaceutical dosage form.

10 11. The pharmaceutical composition of Claim 5 wherein the composition is formulated in an enterically coated time controlled-release pharmaceutical dosage form.

12. The pharmaceutical composition of Claim 1 wherein the composition is formulated in a drug delivery device comprised of:

- 15 (A) a compressed core prepared from an admixture comprising:
- (i) a therapeutically effective amount of the compound;
 - and
 - (ii) a polymer which upon hydration forms gelatinous microscopic particles; and
-
- 20 (B) a water insoluble, water impermeable polymeric coating comprising a polymer and a plasticizer, which surrounds and adheres to the core, the coating having a plurality of formed apertures exposing between about 1 and about 75% of the core surface;
- 25 and wherein the release rate of the compound from the device is a function of the number and size of the apertures.

13. The pharmaceutical composition of Claim 12 wherein the drug delivery device is additionally comprised of an enteric overcoat exterior to the water insoluble, water impermeable polymeric coating.

30

17. The composition of Claim 1 wherein the compound is a pharmaceutically acceptable salt of dihydroxy open acid simvastatin.

5

18. The composition of Claim 1 wherein the compound is a calcium salt of dihydroxy open acid simvastatin.

19. The composition of claim 1 wherein the compound is an ammonium salt of dihydroxy open acid simvastatin.

10

20. The composition of Claim 1 wherein the compound is a crystalline hydrated calcium salt of dihydroxy open acid simvastatin having the solid-state ^{13}C nuclear magnetic resonance spectrum shown in Figure 4.

15

24. The composition of Claim 5 wherein the compound is a pharmaceutically acceptable salt of simvastatin.

25. The composition of Claim 5 wherein the compound is a calcium salt of dihydroxy open acid simvastatin.

20

26. The composition of Claim 5 wherein the compound is an ammonium salt of dihydroxy open acid simvastatin.

27. The composition of Claim 5 wherein the compound is a crystalline hydrated calcium salt of dihydroxy open acid simvastatin having the solid-state ^{13}C nuclear magnetic resonance spectrum shown in Figure 4.

25

28. The composition of claim 25 further comprising BHA.

30

29. The composition of claim 25 further comprising propyl gallate.

30. The composition of claim 25 further comprising BHA and propyl gallate.

31. An oral pharmaceutical composition made by combining a therapeutically effective amount of a compound selected from dihydroxy open acid simvastatin and the pharmaceutically acceptable salts and esters thereof with a pharmaceutically acceptable carrier in a delayed-release dosage form.

32. A process for preparing an oral pharmaceutical composition comprising combining a compound selected from dihydroxy open acid simvastatin and the pharmaceutically acceptable salts and esters thereof with a pharmaceutically acceptable carrier in a delayed-release dosage form.

33. The use of a compound selected from dihydroxy open acid simvastatin and the pharmaceutically acceptable salts and esters thereof for the preparation of an oral medicament adapted for delayed-release wherein substantial release of the compound after oral administration to a patient does not occur until after passage of the medicament through the stomach.

34. The use of Claim 33 wherein at most 10% by weight of the compound is released into the stomach of the patient after administration.

20

35. The use of Claim 33 wherein at most 5% by weight of the compound is released into the stomach of the patient after administration.

36. The use of Claim 33 wherein at most 1% by weight of the compound is released into the stomach of the patient after administration.

25

38. The use of a compound selected from dihydroxy open acid simvastatin and the pharmaceutically acceptable salts and esters thereof for the preparation of an oral medicament adapted for delayed-release useful for delivering at least 90% by weight of the compound in its dihydroxy open acid form to the intestinal mucosa of a patient after oral administration.

30

39. The use of Claim 38 wherein at least 95% by weight of the compound is delivered in its dihydroxy open acid form to the intestinal mucosa of the patient after oral administration.

5 40. The use of Claim 38 wherein at least 99% by weight of the compound is delivered in its dihydroxy open acid form to the intestinal mucosa of the patient after oral administration.

10 42. The use of a compound selected from dihydroxy open acid simvastatin and the pharmaceutically acceptable salts and esters thereof for the preparation of an oral medicament adapted for delayed-release useful for inhibiting HMG-CoA reductase in a patient in need of such treatment wherein substantial release of the compound after oral administration is delayed until after passage of the medicament through the stomach.

15 43. The use of a compound selected from dihydroxy open acid simvastatin and the pharmaceutically acceptable salts and esters thereof for the preparation of an oral medicament adapted for delayed-release useful for treating hypercholesterolemia in a patient in need of such treatment wherein substantial
20 release of the compound after oral administration is delayed until after passage of the medicament through the stomach.

25 44. The use of a compound selected from dihydroxy open acid simvastatin and the pharmaceutically acceptable salts and esters thereof for the preparation of an oral medicament adapted for delayed-release useful for preventing or reducing the risk of developing atherosclerotic disease in a patient at risk of developing atherosclerotic disease wherein substantial release of the compound after oral administration is delayed until after passage of the medicament through the stomach.

30 45. The use of Claim 44 wherein the medicament is useful for preventing or reducing the risk of developing atherosclerotic disease selected from cardiovascular disease, cerebrovascular disease and peripheral vessel disease.

46. The use of Claim 44 wherein the medicament is useful for preventing or reducing the risk of developing coronary heart disease.

5 47. The use of a compound selected from dihydroxy open acid simvastatin and the pharmaceutically acceptable salts and esters thereof for the preparation of an oral medicament adapted for delayed-release useful for treating atherosclerotic disease in a patient in need of such treatment wherein substantial release of the compound after oral administration is delayed until after passage of the medicament through the stomach.

10

48. The use of Claim 47 wherein the medicament is useful for treating atherosclerotic disease selected from cardiovascular disease, cerebrovascular disease and peripheral vessel disease.

15

49. The use of Claim 47 wherein the medicament is useful for treating coronary heart disease.

50. The use of a compound selected from dihydroxy open acid simvastatin and the pharmaceutically acceptable salts and esters thereof for the
20 preparation of an oral medicament adapted for delayed-release useful preventing or reducing the risk of occurrence or recurrence of an atherosclerotic disease event in a person at risk of having an atherosclerotic disease event wherein substantial release of the compound after oral administration is delayed until after passage of the medicament through the stomach.

25

51. The use of Claim 50 wherein the medicament is useful for preventing or reducing the risk of occurrence or recurrence of an atherosclerotic disease event in a person who has atherosclerotic disease.

30

52. The use of Claim 50 wherein the medicament is useful for preventing or reducing the risk of occurrence or recurrence of an atherosclerotic disease event in a person who is at risk of developing atherosclerotic disease.

53. The use of Claim 50 wherein the medicament is useful for preventing or reducing the risk of occurrence or recurrence of an atherosclerotic disease event selected from a coronary heart disease event, a cerebrovascular event and intermittent claudication.

5

54. The use of Claim 50 wherein the medicament is useful for preventing or reducing the risk of occurrence or recurrence of a coronary heart disease event selected from coronary heart disease death, myocardial infarction and coronary revascularization procedures.

10

55. The use of Claim 50 wherein the medicament is useful for preventing or reducing the risk of occurrence or recurrence of a cerebrovascular event selected from a cerebrovascular accident and a transient ischemic attack.

15

58. The use of any of Claims 42 to 55 wherein the compound is a pharmaceutically acceptable salt of dihydroxy open acid simvastatin.

59. The use of any of Claims 42 to 55 wherein the compound is a calcium-salt-of-dihydroxy-open-acid-simvastatin.

20

60. The use of any of Claims 42 to 55 wherein the compound is an ammonium salt of dihydroxy open acid simvastatin.

61. The use of any of Claims 42 to 55 wherein the compound is a crystalline hydrated calcium salt of dihydroxy open acid simvastatin having the solid-state ¹³C nuclear magnetic resonance spectrum shown in Figure 4.

25

62. The composition of Claim 1 wherein the geometric mean ration of the plasma AUC of active vs. total HMG-CoA reductase inhibitory activity is greater than or equal to 90% in the patient after oral administration.

30

63. The composition of Claim 1 wherein the geometric mean ratio of the plasma AUC of active vs. total MHG-CoA reductase inhibitory activity is greater than or equal to 95% in the patient after oral administration.

64. The composition of Claim 1 wherein the geometric mean ratio of the plasma AUC of active vs. total HMG-CoA reductase inhibitory activity is greater than or equal to 99% in the patient after oral administration.

5

STATEMENT UNDER ARTICLE 19 (1)

Claims 1, 31-33, 38, 42-44, 47 and 50 are amended to recite -- dihydroxy open acid simvastatin -- in place of "dihydroxy open acid statin". Claims 2-4, 7, 9, 17-20, 24-27, 34-36, 39-40 and 58-61 are amended to replace the term "statin" with -- compound -- so as to maintain proper and consistent antecedent basis with the claims upon which they depend. In claims 2-4 the word -- oral -- is inserted prior to "administration". In Claims 20, 27 and 61, the XRPD pattern has been replaced with a solid state ^{13}C NMR spectrum. The dependencies of Claims 28-30 are amended from Claim 27 to Claim 25.




FIG.1




FIG.2




FIG.3




FIG. 4

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/02626

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K31/22 A61P3/06 A61K9/28

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5 225 202 A (HODGES GARRY R ET AL) 6 July 1993 (1993-07-06) the whole document	1-61
X	EP 0 547 000 A (SANDOZ LTD) 16 June 1993 (1993-06-16) page 2, line 3,4,35-40 page 10; examples 7-9	1-61
X,P	WO 99 30692 A (ANDRX PHARMACEUTICALS INC) 24 June 1999 (1999-06-24) examples 1-4	1-61
A	WO 97 23200 A (KOWA CO ;NISSAN CHEMICAL IND LTD (JP); MURAMATSU TOYOJIRO (JP); MA) 3 July 1997 (1997-07-03) the whole document	1-61

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

30 May 2000

Date of mailing of the international search report

09/06/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Trifilieff-Riolo, S

INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No

PCT/US 00/02626

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 5225202 A	06-07-1993	NONE	
EP 0547000 A	16-06-1993	AT 401872 B	27-12-1996
		AT 190595 A	15-05-1996
		AT 401870 B	27-12-1996
		AT 244992 A	15-05-1996
		AU 661075 B	13-07-1995
		AU 3006992 A	17-06-1993
		CA 2085037 A	13-06-1993
		CH 684309 A	31-08-1994
		CZ 9203633 A	15-09-1993
		CY 1994 A	05-09-1997
		DE 4240430 A	17-06-1993
		FI 925615 A	13-06-1993
		FR 2684876 A	18-06-1993
		GB 2262229 A,B	16-06-1993
		HK 25597 A	06-03-1997
		HU 63328 A	30-08-1993
		IL 104041 A	27-12-1998
		IT 1256698 B	12-12-1995
		JP 2774037 B	09-07-1998
		JP 5246844 A	24-09-1993
		LU 88201 A	09-09-1994
		MX 9207152 A	01-07-1993
		NO 302099 B	26-01-1998
		NZ 245421 A	27-11-1995
		NZ 270729 A	27-11-1995
		RO 111542 A	29-11-1996
		SK 363392 A	09-11-1994
		US 5356896 A	18-10-1994
		ZA 9209642 A	13-06-1994
WO 9930692 A	24-06-1999	US 5916595 A	29-06-1999
		AU 1627599 A	05-07-1999
WO 9723200 A	03-07-1997	AU 1171597 A	17-07-1997
		CA 2213608 A	03-07-1997
		CZ 9702681 A	13-05-1998
		EP 0814782 A	07-01-1998
		JP 11503763 T	30-03-1999
		NO 973814 A	13-10-1997
		NZ 324446 A	30-08-1999
		PL 321868 A	22-12-1997
		SK 116097 A	04-03-1998

1/9/1

012141471

WPI Acc No: 1998-558383/199848

XRAM Acc No: C98-167255

Use of hypocholesterolaemic agents to prevent or treat diseases
involving prion conformation changes e.g. scrapie, BSE, Alzheimer's disease
and CJD - by reducing number of shingolipid-cholesterol rafts in plasma
membranes

Patent Assignee: EMBL EURO LAB MOLEKULARBIOLOGIE (EMBL-N)

Inventor: SIMONS K

Number of Countries: 020 Number of Patents: 002

Patent Family:

Patent No	Kind	Date	Applicat No	Kind	Date	Week
DE 19716120	A1	19981022	DE 1016120	A	19970417	199848 B
WO 9847518	A1	19981029	WO 98EP2284	A	19980417	199849

Priority Applications (No Type Date): DE 1016120 A 19970417

Patent Details:

Patent No	Kind	Lan	Pg	Main IPC	Filing Notes
-----------	------	-----	----	----------	--------------

DE 19716120	A1		4	A61K-038/17	
-------------	----	--	---	-------------	--

WO 9847518	A1	G		A61K-031/715	
------------	----	---	--	--------------	--

Designated States (National): JP US

Designated States (Regional): AT BE CH CY DE DK ES FI FR GB GR IE IT LU MC
NL PT SE

Abstract (Basic): DE 19716120 A

Use of hypocholesterolaemic agents (I) for prophylaxis or treatment of
diseases dependent on conformation changes in prions or of Alzheimer's
disease and for controlling signal transduction processes in cell
membranes, is new.

USE - (I) reduce the number of sphingolipid-cholesterol rafts in
plasma membranes, thus reducing the number of possible anchoring points at
which conformational changes of proteins can occur. Such rafts also
contain numerous proteins which play an important role in signal
transduction. (I) are useful for treating scrapie in sheep, BSE in cattle,
Alzheimer's disease and related disorders e.g. Creutzfeldt-Jakob disease.

ADVANTAGE - No therapy has previously been available for these
diseases.

Dwg.0/1

Title Terms: HYPOCHOLESTEROLAEMIC; AGENT; PREVENT; TREAT; DISEASE;
CONFORMATION; CHANGE; DISEASE; REDUCE; NUMBER; CHOLESTEROL; RAFT; PLASMA;
MEMBRANE

Derwent Class: B05; C03

International Patent Class (Main): A61K-031/715; A61K-038/17

File Segment: CPI

Manual Codes (CPI/A-N): B14-J01A4; B14-N16; B14-J01A4; C14-J01A4; B14-N16;

C14-N16; C14-J01A4; C14-N16

Chemical Fragment Codes (M1):

01 M423 M781 M903 P446 P625 V797

Derwent WPI (Dialog® File 351): (c) 2001 Derwent Info Ltd. All rights reserved.

© 2001 The Dialog Corporation

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☒ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.